

VI. BIBLIOGRAFIA

Abraham K, Knoll A, Ende M, Papke O, Helge H (1996). Intake, fecal excretion, and body burden of polychlorinated dibenzo-p-dioxins and dibenzofurans in breast-fed and formula-fed infants. *Pediatr Res* 40: 671-679.

Alcock RE, Gemmill R, Jones KC (1999). Improvements to the UK PCDD/F and PCB atmospheric emission inventory following an emission measurement programme. *Chemosphere* 38: 759-770.

Amaral Mendes JJ (2002). The endocrine disrupters: a major medical challenge. *Food Chem Toxicol* 40:781-788.

Aoki Y (2001). Polychlorinated biphenyls, polychlorinated dibenzo-p-dioxins, and polychlorinated dibenzofurans as endocrine disrupters- What we have learned from Yusho disease. *Environ Res* 86: 2-11.

Arija V, Salas J, Fernández J, Cucó G, Martí C (1996). Consumo, hábitos alimentarios y estado nutricional de la poblacions de Reus (IX). Evolución del consumo de alimentos, de su participación en la ingestión de energía y nutrientes y de su relación con el nivel socioeconómico y cultural entre 1983 y 1993. *Med Clin* 106: 174-179.

ATSDR (2000). Polychlorinated biphenyls. Agency for Toxic Substances and Disease Registry, Atlanta, GA. <http://www.atsdr.cdc.gov/tfacts17.html>.

ATSDR (2002). Polybrominated Biphenyls and Polybrominated Diphenyl Ethers (PBBs and PBDEs). <http://www.atsdr.cdc.gov/toxprofiles/tp68.html>.

Avkhimenko M (2000). Medical and ecological consequences of environmental contamination with PCBs, in: Polychlorinated Biphenyls-Supertoxicants of XXI Century, VINITI Pres, Moscow: 14-30.

Baker JI, Hites RA (2000). Is combustion the major source of polychlorinated dibenz-p-dioxins and dibenzofurans to the environment? A mass balance investigation. Environ Sci Technol 34: 2879-2886.

Bascompta O, Montaña MJ, Martí R, Broto-Puig F, Comellas L, Díaz-Ferrero J, Rodriguez-Larena MC (2002). Levels of persistent organic pollutants (PCDD/F and dioxin-like PCB) in food from the Mediterranean diet. Organohalogen Compd 57: 149-151.

Becker M, Phillips T, Safe S (1991). Polychlorinated diphenyl ethers: a review. Toxicol Environ Chem 33: 189-200.

Birnbaum LS (1994). The mechanism of dioxin toxicity: relationship to risk assessment. Environ Health Perspect 102: 157-167.

Bocio A, Domingo JL (2004) Daily intake of polychlorinated dibenzo-p-dioxins/polychlorinated dibenzofurans (PCDD/PCDFs) in foodstuffs consumed in Tarragona, Spain: a review of recent studies (2001–2003) on human PCDD/PCDF exposure through the diet. Environ Res: *en premsa*.

Bordado J, Ferreira H, Gomes J (1999). Dioxinas e dibenzofuranos no meio ambiente. Departamento de Engenharia Química. Instituto Superior Técnico. Av. Rovisco Pais, 1906 Lisboa Codex.

Branchi I, Capone F, Alleva E, Costa LG (2003). Polybrominated diphenyl ethers: neurobehavioral effects following developmental exposure. Neurotoxicology 24:449-462.

Buekens A, Stieglitz L, Hell K, Huang H, Segers P (2001). Dioxins from thermal and metallurgical processes: recent studies for the iron and steel industry. Chemosphere 42: 729-735.

Capdevila F, Llop D, Guillén N, Luque V, Perez S, Selles V, Fernandez-Ballart J, Martí-Henneberg C (2000). Consumo, hábitos alimentarios y estado nutricional de la población de Reus (X). Evolución de la ingesta alimentaria y de la contribución de los macronutrientes al aporte energético (1983-1999). Med Clin 115, 7-14.

CAPV (Comunidad Autónoma del País Vasco) (1997) Vigilancia de la contaminación química de los alimentos en la Comunidad Autónoma del País Vasco, 1990-1995. Servicio Central de Publicaciones del Gobierno Vasco.

Casarett i Doull's (2001) Toxicology. The basic science of poisons. McGraw Hill, 6th edició.

Chen HL, Lee CC, Liao PC, Guo YL, Chen CH, Su HJ (2003). Associations between dietary intake and serum polychlorinated dibenzo-p-dioxin and dibenzofuran (PCDD/F) levels in Taiwanese. Environ Res 91: 172-178.

Chiarenzelli JR, Alexander C, Isley A, Scrudato R, Pagano J, Ramirez W (2001). Polychlorinated biphenyls in nonaccumulating, century-old sediments: sources, signatures, and mechanism of introduction. Environ Sci Technol 35: 2903-2908.

Choi JW, Fujimaki S, Kitamura K, Hashimoto S, Ito H, Suzuki N Ito H, Suzuki N, Sakai S, Morita M (2002). Polybrominated dibenzo-p-dioxins (PBDDs), dibenzofurans (PBDFs) and diphenyl ethers (PBDEs) in Japanese human adipose tissue. Organohalog Compd 58: 169-171.

Chu S, Covaci A, Schepens P (2003). Levels and chiral signatures of persistent organochlorine pollutants in human tissues from Belgium. Environ Res 93: 167-176.

Coleman PJ, Lee RGM, Alcock RE, Jones KC (1997). Observations on PAH, PCB, and PCDF/F trends in UK urban air, 1991-1995. Environ Sci Technol 31: 2120-2124.

Cuervo L, Jalón M., Rose M, Fernandes A, White S, González de Galdeano L (2002). Dietary intakes of PCDDs, PCDFs and PCBs in total diet samples from the Basque Country (Spain). Organohalogen Compd 55: 219-222.

Darnerud PO (2003). Toxic effects of brominated flame retardants in man and in wildlife. Environ Int 29: 841-853.

Darnerud PO, Eriksen GS, Jóhannesson T, Larsen PB, Viluksela M (2001). Polybrominated diphenyl ethers: occurrence, dietary exposure, and toxicology. Environ Health Perspect 109: 49-68.

De Boer J, Denneman M (1998). Polychlorinated diphenylethers: origin, analysis, distribution, and toxicity in the marine environment. Rev Environ Contam Toxicol 157: 131-144.

De Wit (2002). An overview of brominated flame retardants in the environment. Chemosphere 46: 583-624.

DOGC (1986). Ordre de 9 de setembre de 1986, de limitació de l'ús dels policlorobifenils i els policloroterfenils. Departament d'Indústria, Comerç i Turisme.

Domingo JL, Schuhmacher M, Granero S, Llobet JM (1999). PCDDs and PCDFs in food samples from Catalonia, Spain. As assessment of dietary intake. Chemosphere 38: 3517-3528.

Domingo JL, Agramunt MC, Nadal M, Schuhmacher M, Corbella J (2002). Health risk assessment of PCDD/PCDF exposure for the population living in the vicinity of a municipal waste incinerator. Arch Environ Contam Toxicol 43: 461-465.

Eljarrat E, Monjonell A, Caixach J, Rivera J (2002). Toxic potency of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and polychlorinated biphenyls in food samples from Catalonia (Spain). *J Agric Food Chem* 50: 1161-1167.

European Commission Summary Record of the 83rd Meeting of the Standing Committee on Foodstuffs, July 19-20, 2001, Brussels, Belgium.

Fiedler H (1996). Sources of PCDD/PCDF and impact on the environment. *Chemosphere* 32: 55-64.

Fiedler H (1998). Thermal formation of PCDD/PCDF: a survey. *Environ Eng Sci* 15: 49-58.

Fiedler H (1999). Dioxin and furan inventories. National and Regional Emissions of PCDD/PCDF. UNED chemicals, Geneva, Switzerland.

Fiedler H, Hutzinger O (1990). Dioxins: sources of environmental load and human exposure. *Toxicol Environ Chem* 29: 157-234.

Flesch-Janys D, Becher H, Gurn P, Jung D, Konietzko J, Manz A, Papke O (1996). Elimination of polychlorinated dibenzo-p-dioxins and dibenzofurans in occupationally exposed persons. *J Toxicol Environ Health* 47: 363-378.

Focant JF, Eppe G, Pirard C, Massart AC, André JE, De Pauw E (2002). Levels and congener distributions of PCDDs, PCDFs and non-*ortho* PCBs in Belgian foodstuffs. Assessment of dietary intake. *Chemosphere* 48: 167-179.

Fries GF (1995). A review of the significance of animal food products as potential pathways of human exposures to dioxins. *J Anim Sci* 73: 1639-1650.

Gilbert ME (2003). Perinatal exposure to Polychlorinated Biphenyls alters excitatory synaptic transmission and short-term plasticity in the hippocampus of the adult rat. *Neurotoxicology* 24: 851-860.

Gilman H, Dietrich J (1957). *J Amer Chem Soc* 79: 1439. En: Soriano V. Dioxinas ([http://www.opolanco.es/Apat/Boletín 14/dioxinas.htm](http://www.opolanco.es/Apat/Boletín%2014/dioxinas.htm)).

Guvenius DM, Aronsson A, Ekman-Ordeberg G, Bergman A, Noren K (2003). Human prenatal and postnatal exposure to polybrominated diphenyl ethers, polychlorinated biphenyls, polychlorobiphenylols, and pentachlorophenol. *Environ Health Perspect* 111: 1235-1241.

Hagmar L, Sjödin A, Höglund P, Thuresson K, Rylander L, Bergman A (2000). Biological half-lives of polybrominated diphenyl ethers and tetrabromobisphenol A in exposed workers. *Organohalog Compd* 47: 198-201.

Hakk H, Letcher RJ (2003). Metabolism in the toxicokinetics and fate of brominated flame retardants: a review. *Environ Int* 29: 801-828.

Harrad SJ, Jones KC (1992). A source inventory and budget for chlorinated dioxins and furans in the United Kingdom environment. *Sci Total Environ* 126: 89-107.

Hays SM, Aylward LL (2003). Dioxin risks in perspective: past, present, and future. *Regul Toxicol Pharmacol* 37: 202-217.

Hites RA, Foran JA, Carpenter DO, Hamilton MC, Knuth BA, Schwager SJ (2004). Global assessment of organic contaminants in farmed salmon. *Science* 303: 226-229.

Hsu MS, Cheng PS, Ma E, Chou U, Chen LP, Jone CH, Chou SS, Cheng CC, Yu CY, Liao CH, Ling YC (2002). A preliminary total diet study on PCDD/Fs-intake from food in Taiwan. *Organohalogen Compd* 55: 231-234.

Hutzinger O, Fielder H (1989). Sources and emissions of PCDD/PCDF. Chemosphere 18: 23-32.

IARC (1990). Monographs: some flame retardants and textile chemicals, and exposure in the textile manufacturing industry. Decabromodiphenyl Oxide, vol. 48: 73-84. France: Lyon.

IARC (1995). Monograph Volumes 1972-1994, WHO, Geneva, Switzerland, pp 1-60, <http://www.holgr/medical/nuclear/carcinogen.htm>.

Ikonomou MG, Rayne S, Addison RF (2002). Exponential increases of the brominated flame retardants, polybrominated diphenyl ethers, in the Canadian Arctic from 1981 to 2000. Environ Sci Technol 36: 1886-1892.

Iverson F, Newsome H, Hierlihy L (1987). Induction of rat hepatic monooxygenase activity by polychlorinated diphenyl ethers. Chem Toxic 25: 305-307.

Kim JG, Kim KS, Joo CH, You JC (2000). Exposure of PCDD/Fs via air and food in Koreans. Organohalogen Compd 47: 314-318.

Kimbrough RD (1995). Polychlorinated biphenyls (PCBs) and human health: An update. Crit Rev Toxicol 25:133-163.

Kitamura K, Kikichi Y, Watanabe S (2000). Health effects of chronic exposure to polychlorinated dibenzo-p-dioxins (PCDD), dibenzofurans (PCDF) and coplanar PCB (Co-PCB) of municipal waste incinerator workers. J Epidemiol 10: 262-270.

Kiviranta H, Hallikainen A, Ovaskaiens ML, Kumpulainen J, Vartiainen T (2001). Dietary intakes of polychlorinated dibenzo-p-dioxins, dibenzofurans and polychlorinated biphenyls in Finland. Food Add Contam 18: 945-953.

Knoth W, Mann W, Meyer R, Nebhuth J (2002). Polybrominated diphenylether in house dust. *Organohalog Compd* 58:213-216.

Kogevinas M, Janer G (2000). Health effects of dioxins. *Med Clin* 115: 740-748.

Kogevinas M (2001). Human health effects of dioxins: cancer, reproductive and endocrine system effects. *Hum Reprod Update* 7: 331-339.

Koistinen J (2000). Polychlorinated Diphenyl Ethers (PCDE). In: *Handbook of Environmental Chemistry Vol 3, Part K, Chapter 7* (Paasivirta J, ed) Berlin/Heidelberg. Germany: Springer-Verlag.

Koistinen J, Mussalo-Rauhamaa H, Paasivirta J (1995). Polychlorinated diphenyl ethers, dibenzo-p-dioxins and dibenzofurans in Finnish human tissues compared to environmental samples. *Chemosphere* 31: 4259-4271.

Koistinen J, Stenman O, Haahti H, Suonperä M, Paasivirta J (1997). Polychlorinated diphenyl ethers, dibenzo-p-dioxins, dibenzofurans and biphenyls in seals and sediment from the gulf of Finland. *Chemosphere* 35: 1249-1269.

Kreis K (1985). Studies on populations exposed to polychlorinated biphenyls. *Environ Health Perspect* 60:193-199.

Kreuzer PE, Csanady GA, Baur C, Kessler W, Papke O, Greim H, Filser JG (1997). 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and congeners in infants. A toxicokinetic model of human lifetime body burden by TCDD with special emphasis on its uptake by nutrition. *Arch Toxicol* 71: 383-400.

Kutz FW, Barnes DG, Bottimore DP (1990). The international toxicity equivalency factor (I-TEF) method of risk assessment for complex mixtures of dioxins and related compounds. *Chemosphere* 20: 751-757.

Ladron de Guevara J, Moya Pueyo V (1995). Toxicología Médica. Clínica y Laboral. Interamericana, McGraw-Hill, Madrid.

LaKind JS, Filser JG (1999). Dietary exposure to PCBs and dioxins. Environ Health Perspect 107: 495-497.

Lazaro R, Herrera A, Conchello MP, Ariño AA, Bayarri S, Yagüe C, Peiro JM (1999). Levels of selected polychlorinated biphenyl congeners in total diet samples from Aragón, Spain. J Food Prot 9:1054-1058.

Lemesh RA (1992). Polychlorinated biphenyls: An overview of metabolic, toxicologic and health consequences. Vet Hum Toxicol 34: 256-260.

Liem AK, Furst P, Rappe C (2000). Exposure of populations to dioxins and related compounds. Food Addit Contam 17:241-259.

Lind Y, Aune M, Atuma S, Becker W, Bjerselius R, Glynn A, Darnerud PO (2002). Food intake of the brominated flame retardants PBDEs and HBCD in Sweden. Organohalogen Compd 58: 181-184.

Llibre Blanc. Evaluació de l'estat nutricional de la població catalana (1992-93) Generalitat de Catalunya. Departament de Sanitat i Seguretat Social.

MAPA (2000). Alimentación en España. Ministerio de Agricultura, Pesca y Alimentación.

Mastrantonio G (2000). Compuestos organoclorados como contaminantes persistentes: el caso de las dioxinas y los bifenilos policlorados. Progreso SA- Grupo Consultores Independiente. Nota técnica nº2: 1-2.

McDonald TA (2002). A perspective on the potential health risks of PBDEs. Chemosphere 46: 745-755.

McGregor DB, Partensky C, Wilbourn J, Rice JM (1998). An IARC evaluation of polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans as risk factors in human carcinogenesis. Environ Health Perspect 106: 755-760.

Meironyte Guvenius D, Bergman A, Noren K (2001). Polybrominated diphenyl ethers in Swedish human liver and adipose tissue. Arch Environ Contam Toxicol 40: 564-570.

Michalek JE, Tripathi RC, Kulkarni PM, Pirkle JL (1996). The reliability of the serum dioxin measurement in veterans of Operation Ranch Hand. J Expo Anal Environ Epidemiol 6: 327-338.

Negri E, Bosetti C, Fattore E, La Vecchia C (2003) Environmental exposure to polychlorinated biphenyls (PCBs) and breast cancer: a systematic review of the epidemiological evidence. Eur J Cancer Prev 12: 509-516.

Norris JM, Kociba RJ, Schwetz BA, Rose JQ, Humiston CG, Jewett GL, Gehring PJ, Mailhes JB (1975). Toxicology of octabromobiphenyl and decabromodiphenyl oxide. Environ Health Perspect 11: 153-161.

NRC (1994) Science and Judgment in Risk Assessment. Washington, DC: National Academy Press.

OECD (1994a). Diari Oficial de les Comunitats Europees. Posició Comú N° 26/94 relativa a la incineració de residus perillosos, N°C232: 35-50.

OECD (1994b). Selected brominated flame retardants, risk reduction monograph 3. Paris, France: OECD, Environment Directorate.

Omenn GS, Faustman EM (1997). Risk assessment and risk management, in Detels R, Holland W, McEwen J, Omenn G (eds). Oxford Textbook of Public Health, 3d ed. New York. Oxford University Press, pp 969-986. (4th ed. in press.).

Otha S, Ishikuza D, Nishimura H, Nakao T, Aozasa A, Shimidzu Y, Ochiai F, Kida T, Nishi M, Miyata H (2002). Comparison of polybrominated diphenyl ethers in fish, vegetables, and meats and levels in human milk of nursing women in Japan. Chemosphere 46: 689-696.

Parzefall W (2002). Risk assessment of dioxin contamination in human food. Food Chem Toxicol 40: 1185-1189.

Rahman F, Langford KH, Scrimshaw MD, Lester JN (2001). Polybrominated diphenyl ethers (PBDE) flame retardants. Sci Total Environ 275: 1-17.

Rappe C (1992). Sources of PCDDs and PCDFs. Introduction, reactions, levels, patterns, profiles and trends. Chemosphere 25: 41-44.

Rose M, White S, Harrison N, Gleadle A (2001). PCDD/Fs (dioxins) and PCBs in the UK diet: 1997 total diet study samples. 21st International Symposium on Halogenated Environmental Organic Pollutants & POPs. Kyongju, Korea.

Rosiak K, Li MH, Degitz SJ, Skalla DW, Chu I, Francis BM (1997). Maternal and developmental toxicity of polychlorinated diphenyl ethers (PCDEs) in Swiss-Webster mice and Sprague Dawley rats. Toxicology 121: 191-204.

Ryan JK, Patry B (2001). Body burdens and food exposure in Canada for polybrominated diphenyl ethers (BDEs). Organohalogen Compd 51: 226-229.

Rylander L, Strömberg U, Hagmar L (1996). Dietary intake of fish contaminated with persistent organochlorine compounds in relation to low birthweight. *Scand J Work Environ Health* 22: 260-266.

Rylander L, Strömberg U, Dyremark E, Östman C, Nilsson-Ehle P, Hagmar L (1998). Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to low birth weight. *Am J Epidemiol* 147: 493-502.

Safe S (1990). Polychlorinated Biphenyls (PCBs), Dibenz-p-Dioxins (PCDDs), Dibenzofurans (PCDFs), and related compounds: Environmental and mechanistic considerations which support the development of toxic equivalency factors (TEFs). *Crit Rev Toxicol* 21: 51-88.

Safe SH (1994). Polychlorinated biphenyls (PCBs): Environmental impact, biochemical and toxic responses, and implications for risk assessment. *Crit Rev Toxicol* 24:87-149.

Schechter A, Cramer P, Boggess K, Stanley J, Papke O, Olson J, Silver A, Schmitz M (2001). Intake of dioxins and related compounds from food in the US population. *J Toxicol Environ Health* 63: 1-18.

Schmid P, Gujer E, Zennegg M, Studer C (2003). Temporal and local trends of PCDD/F levels in cow's milk in Switzerland. *Chemosphere* 53: 129-136.

Schuhmacher M, Agramunt MC, Rodriguez-Larena MC, Diaz-Ferrero J, Domingo JL (2002). Baseline levels of PCDD/Fs in soil and herbage samples collected in the vicinity of a new hazardous waste incinerator in Catalonia, Spain. *Chemosphere* 46: 1343-1350.

Seta N, Arfi C, Momas I (2000). General population exposure to dioxins, human ubiquitous pollutants. *Rev Epidemiol Sante Publique* 48: 281-293.

Shadel B, Evans NRG, Roberts D, Clardy S, Jordan-Izaguirre SD, Patterson Jr DG, Needham LL (2001). Background levels of non-ortho-substituted (coplanar) polychlorinated biphenyls in human serum of Missouri residents. Chemosphere 43: 967-976.

She J, Petreas M, Winkler J, Visita P, McKinney M, Kopec D (2002). PBDEs in the San Francisco Bay Area: measurements in harbor seal blubber and human breast adipose tissue. Chemosphere. 46: 697-707.

Sjödin A, Hagmar L, Klasson-Wehler E, Kronholm-Diab K, Jakobsson E, Bergman A (1999). Flame retardant exposure: polybrominated diphenyl ethers in blood from Swedish workers. Environ Health Perspect 107: 643-648.

Sjödin A, Patterson Jr D G, Bergman A (2003). A review on human exposure to brominated flame retardants-particularly polybrominated diphenyl ethers. Environ Int 29: 829-839.

Stafford CJ (1983). Halogenated diphenylethers in avian tissues and eggs by GC/MS. Chemosphere 12: 1487-1495.

Steenland K, Deddens J, Piacitelli L (2001). Risk assesment for 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) based on an epidemiologic study. Am J Epidemiol 154: 451-458.

Sweeney A, Symanski E, Burau KD, Kim YJ, Humphrey HEB, Smith MS (2001). Changes in serum PBB and PCB levels over time among women of varying ages at exposure. Environ Res Sect 86: 128-139.

Sweetman AJ, Alcock RE, Wittsiepe J, Jones KC (2000) Human exposure to PCDD/Fs in the UK: the development of a modeling approach to give historical and future perspectives. Environ Int 26: 37-47.

Thoma H, Mucke W, Kauert G (1990). Comparison of the polychlorinated dibenz-p-dioxin and dibenzofurans in human tissue and human liver. Chemosphere 20: 433-442.

Thuresson K, Jakobsson K, Hagmar L, Sjödin A, Bergman A (2002). Decabromodiphenyl ether exposure to workers manufacturing rubber and in an industrial setting producing rubber coated electric wires. Organohalog Comp 58: 165-168.

Toyoda M, Uchibe H, Yanagi T, Kono Y, Hori T, Ida T (1999). Dietary daily intake of PCDDs, PCDFs and coplanar PCBs by total diet study in Japan. J Food Hyg Soc Japan 40: 98-110.

Tsutsumi T, Yanagi T, Nakamura M, Kono Y, Uchibe H, Iida T, Hori T, Nakagawa R, Tobiishi K, Matsuda R, Sasaki K, Toyoda M (2001). Update of daily intake of PCDDs, PCDFs, and dioxin-like PCBs from food in Japan. Chemosphere 45: 1129-1137.

US EPA (1993). Integrated Risk Information Center. Reference Dose (RfD): Description and Use in Health Risk Assessments Background Document 1A March 15.

US EPA (2000). Exposure and human health reassessment of 2,3,7,8-tetrachlorodibenzo-p-dioxins and related compounds. EPA/600/P-00/001 Bg.

US EPA (2003a). Draft final guidelines for carcinogen risk assessment (external review draft, february 2003). other ncea-f-0644a. U.S. Environmental Protection Agency, Risk Assessment Forum, Washington, DC.

US EPA (2003b). Glossary of IRIS Terms. Integrated Risk Information System. Environmental Protection agency. <http://www.epa.gov/iris/gloss8.htm>.

Van Birgelen APJM, van den Berg M (2000). Toxicokinetics. *Food Addit Contam* 17: 267-273.

Van den Berg M, Peterson RE, Schrenk D (2000). Human risk assessment and TEFs. *Food Addit Contam* 17: 347-358.

Van Larebeke N, Hens L, Schepens P, Covaci A, Baeyens J, Everaert K, Bernheim JL, Vlietinck R, De Poorter G (2001). The Belgian PCB and dioxin incident of January-June 1999: exposure data and potential impact on health. *Environ Health Perspect* 109: 265-273.

Van Leeuwen FXR, Feeley M, Schrenk D, Larsen JC, Farland W, Younes M (2000). Dioxins: WHO's tolerable daily intake revisited. *Chemosphere* 40: 1095-1101.

Vieth B, Heinrich-Hirsch B, Mathar W (2000). Trends in dioxin intake and human milk levels in Germany. *Organohalogen Compd* 47: 300-303.

Vikelsoe J, Johansen E (2000). Estimation of dioxin emission from fires in chemicals. *Chemosphere* 40: 165-175.

Villeneuve JY, Niimi AJ, Metcalfe CD (1999). Distribution and bioaccumulation of chlorinated diphenyl ethers in a contaminated embayment of Lake Ontario. *J Great Lakes Res* 25: 760-771.

Walkowiak J, Wiener JA, Fastabend A, Heinzow B, Kramer U, Schmidt E, Steingruber, HJ, Wundram S, Winneke G (2001). Environmental exposure to polychlorinated biphenyls and quality of the home environment: effects on psychodevelopment in early childhood. *Lancet* 358: 1602-1607.

WHO (1962). Principles in governing consumer safety in relation to pesticide residues. *WHO Tech Rep Ser* 240.

WHO (1985). Guidelines for the Study of Dietary Intakes of Chemical Contaminants. World Health Organization Offset Publication No. 87. Geneva.

WHO (1991). Consultation on Tolerable Daily Intake from food of PCDDs and PCDFs. WHO Regional Office for Europe. EUR/ICP/PCS 030.

WHO (1994). Environmental Health Criteria 162. Brominated diphenyl ethers. Geneva, Switzerland: International Programme on Chemical Safety.

WHO (1997). Environmental Health Criteria 162. Flame retardants: a general introduction. Geneva, Switzerland: International Programme on Chemical Safety.

Wijesekera R, Halliwell C, Hunter S, Harrad S (2002). A preliminary assessment of UK human exposure to polybrominated diphenyl ethers (PBDEs). *Organohalogen Compd* 55: 239-242.

Williams DT, LeBel GL (1988). Chlorinated diphenyl ethers in human adipose tissue *Chemosphere* 17: 2349-2354.

Williams DT, Kennedy B, LeBel GL (1991). Chlorinated diphenyl ethers in human adipose tissue. Part 2. *Chemosphere* 23: 601-608.

Wu WZ, Li W, Xu Y, Wang JW (2001). Long-term toxic impact of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the reproduction, sexual differentiation, and development of different life stages of *Gobiocypris rarus* and *Daphnia magna*. *Ecotoxi & Environ Safety* 48: 293-300.

Wu Y, Li J, Zhao Y, Chen Z, Li W, Chen J (2002). Dietary intake of polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) in populations from China. *Organohalogen Compd* 57: 221-223.

Yoshimura T (2003). Yusho in Japan. Ind Health 41:139-148.