

VII. BIBLIOGRAFIA

A

- Abbracchio MP and Burnstock G (1998) Purinergic signalling: pathophysiological roles. *Jpn J Pharmacol* **78**:113-45.
- Ackley MA, Governo RJ, Cass CE, Young JD, Baldwin SA and King AE (2003) Control of glutamatergic neurotransmission in the rat spinal dorsal horn by the nucleoside transporter ENT1. *J Physiol* **548**:507-17.
- Agarwal RP and Olivero OA (1997) Genotoxicity and mitochondrial damage in human lymphocytic cells chronically exposed to 3'-azido-2',3'-dideoxythymidine. *Mutat Res* **390**:223-31.
- Anderson CM, Xiong W, Young JD, Cass CE and Parkinson FE (1996) Demonstration of the existence of mRNAs encoding N1/cif and N2/cit sodium/nucleoside cotransporters in rat brain. *Brain Res Mol Brain Res* **42**:358-61.
- Andersson U, Filipsson K, Abbott CR, Woods A, Smith K, Bloom SR, Carling D and Small CJ (2004) AMP-activated protein kinase plays a role in the control of food intake. *J Biol Chem* **279**:12005-8.
- Avery TL, Rehg JE, Lumm WC, Harwood FC, Santana VM and Blakley RL (1989) Biochemical pharmacology of 2-chlorodeoxyadenosine in malignant human hematopoietic cell lines and therapeutic effects of 2-bromodeoxyadenosine in drug combinations in mice. *Cancer Res* **49**:4972-8.

B

- Baas AF, Kuipers J, van der Wel NN, Batlle E, Koerten HK, Peters PJ and Clevers HC (2004a) Complete polarization of single intestinal epithelial cells upon activation of LKB1 by STRAD. *Cell* **116**:457-66.
- Baas AF, Smit L and Clevers H (2004b) LKB1 tumor suppressor protein: PARtaker in cell polarity. *Trends Cell Biol* **14**:312-9.
- Baldwin SA, Beal PR, Yao SY, King AE, Cass CE and Young JD (2004) The equilibrative nucleoside transporter family, SLC29. *Pflugers Arch* **447**:735-43.
- Baldwin SA, Mackey JR, Cass CE and Young JD (1999) Nucleoside transporters: molecular biology and implications for therapeutic development. *Mol Med Today* **5**:216-24.
- Baldwin SA, Yao SY, Hyde RJ, Ng AM, Foppolo S, Barnes K, Ritzel MW, Cass CE and Young JD (2005) Functional characterization of novel human and mouse equilibrative nucleoside transporters (hENT3 and mENT3) located in intracellular membranes. *J Biol Chem* **280**:15880-7.
- Barnes K, Ingram JC, Porras OH, Barros LF, Hudson ER, Fryer LG, Foufelle F, Carling D, Hardie DG and Baldwin SA (2002) Activation of GLUT1 by metabolic and osmotic stress: potential involvement of AMP-activated protein kinase (AMPK). *J Cell Sci* **115**:2433-42.
- Barthel A, Schmoll D, Kruger KD, Roth RA and Joost HG (2002) Regulation of the forkhead transcription factor FKHR (FOXO1a) by glucose starvation and AICAR, an activator of AMP-activated protein kinase. *Endocrinology* **143**:3183-6.
- Bartlett GR and Bucolo G (1968) The metabolism of ribonucleoside by the human erythrocyte. *Biochim Biophys Acta* **156**:240-53.
- Becker K, Stegenga S and Conway S (1995) Role of insulin-like growth factor I in regulating growth hormone release and feedback in the male rat. *Neuroendocrinology* **61**:573-83.
- Belt JA, Marina NM, Phelps DA and Crawford CR (1993) Nucleoside transport in normal and neoplastic cells. *Adv Enzyme Regul* **33**:235-52.
- Bergeron R, Russell RR, 3rd, Young LH, Ren JM, Marcucci M, Lee A and Shulman GI (1999) Effect of AMPK activation on muscle glucose metabolism in conscious rats. *Am J Physiol* **276**:E938-44.
- Blake HH and Henning SJ (1983) Weaning in the rat: a study of hormonal influences. *Am J Physiol* **244**:R537-43.
- Bode AM and Dong Z (2002) The paradox of arsenic: molecular mechanisms of cell transformation and chemotherapeutic effects. *Crit Rev Oncol Hematol* **42**:5-24.
- Bontemps F, Mimouni M and Van den Berghe G (1993) Phosphorylation of adenosine in anoxic hepatocytes by an exchange reaction catalysed by adenosine kinase. *Biochem J* **290** (Pt 3):679-84.

- Bouffard DY and Momparler RL (1995) Comparison of the induction of apoptosis in human leukemic cell lines by 2',2'-difluorodeoxycytidine (gemcitabine) and cytosine arabinoside. *Leuk Res* **19**:849-56.
- Boyle DL, Kowaluk EA, Jarvis MF, Lee CH, Bhagwat SS, Williams M and Firestein GS (2001) Anti-inflammatory effects of ABT-702, a novel non-nucleoside adenosine kinase inhibitor, in rat adjuvant arthritis. *J Pharmacol Exp Ther* **296**:495-500.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**:248-54.
- Brasitus TA and Schachter D (1984) Lipid composition and fluidity of rat enterocyte basolateral membranes. Regional differences. *Biochim Biophys Acta* **774**:138-46.
- Buchheimer RE and Linden J (2004) Purinergic regulation of epithelial transport. *J Physiol* **555**:311-21.
- Bzowska A, Kulikowska E and Shugar D (2000) Purine nucleoside phosphorylases: properties, functions, and clinical aspects. *Pharmacol Ther* **88**:349-425.

C

- Cai J and Lee CW (1996) Tamoxifen inhibits nitrobenzylthioinosine-sensitive equilibrative uridine transport in human MCF-7 breast cancer cells. *Biochem J* **320 (Pt 3)**:991-5.
- Cano-Soldado P, Lorryoz IM, Molina-Arcas M, Casado FJ, Martinez-Picado J, Lostao MP and Pastor-Anglada M (2004) Interaction of nucleoside inhibitors of HIV-1 reverse transcriptase with the concentrative nucleoside transporter-1 (SLC28A1). *Antivir Ther* **9**:993-1002.
- Cao D, Leffert JJ, McCabe J, Kim B and Pizzorno G (2005) Abnormalities in uridine homeostatic regulation and pyrimidine nucleotide metabolism as a consequence of the deletion of the uridine phosphorylase gene. *J Biol Chem* **280**:21169-75.
- Carabaza A, Ricart MD, Mor A, Guinovart JJ and Ciudad CJ (1990) Role of AMP on the activation of glycogen synthase and phosphorylase by adenosine, fructose, and glutamine in rat hepatocytes. *J Biol Chem* **265**:2724-32.
- Carling D (2004) The AMP-activated protein kinase cascade--a unifying system for energy control. *Trends Biochem Sci* **29**:18-24.
- Carling D (2005) AMP-activated protein kinase: balancing the scales. *Biochimie* **87**:87-91.
- Carpenter G (1979) Regulation of epidermal growth factor (EGF) receptor activity during the modulation of protein synthesis. *J Cell Physiol* **99**:101-6.
- Chabes AL, Bjorklund S and Thelander L (2004) S Phase-specific transcription of the mouse ribonucleotide reductase R2 gene requires both a proximal repressive E2F-binding site and an upstream promoter activating region. *J Biol Chem* **279**:10796-807.
- Chandrasena G, Giltay R, Patil SD, Bakken A and Unadkat JD (1997) Functional expression of human intestinal Na⁺-dependent and Na⁺-independent nucleoside transporters in *Xenopus laevis* oocytes. *Biochem Pharmacol* **53**:1909-18.
- Chaudary N, Naydenova Z, Shuralyova I and Coe IR (2004) The adenosine transporter, mENT1, is a target for adenosine receptor signaling and protein kinase C epsilon in hypoxic and pharmacological preconditioning in the mouse cardiomyocyte cell line, HL-1. *J Pharmacol Exp Ther* **310**:1190-8.
- Chaudary N, Shuralyova I, Liron T, Sweeney G and Coe IR (2002) Transport characteristics of HL-1 cells: a new model for the study of adenosine physiology in cardiomyocytes. *Biochem Cell Biol* **80**:655-65.
- Che M, Gatmaitan Z and Arias IM (1997) Ectonucleotidases, purine nucleoside transporter, and function of the bile canalicular plasma membrane of the hepatocyte. *Faseb J* **11**:101-8.
- Che M, Ortiz DF and Arias IM (1995) Primary structure and functional expression of a cDNA encoding the bile canalicular, purine-specific Na⁽⁺⁾-nucleoside cotransporter. *J Biol Chem* **270**:13596-9.
- Cheeseman CI and Harley B (1991) Adaptation of glucose transport across rat enterocyte basolateral membrane in response to altered dietary carbohydrate intake. *J Physiol* **437**:563-75.
- Cheng SW, Fryer LG, Carling D and Shepherd PR (2004) Thr2446 is a novel mammalian target of rapamycin (mTOR) phosphorylation site regulated by nutrient status. *J Biol Chem* **279**:15719-22.

- Chimploy K and Mathews CK (2001) Mouse ribonucleotide reductase control: influence of substrate binding upon interactions with allosteric effectors. *J Biol Chem* **276**:7093-100.
- Choi DS, Cascini MG, Mailliard W, Young H, Paredes P, McMahon T, Diamond I, Bonci A and Messing RO (2004) The type 1 equilibrative nucleoside transporter regulates ethanol intoxication and preference. *Nat Neurosci* **7**:855-61.
- Christofi FL, Zhang H, Yu JG, Guzman J, Xue J, Kim M, Wang YZ and Cooke HJ (2001) Differential gene expression of adenosine A1, A2a, A2b, and A3 receptors in the human enteric nervous system. *J Comp Neurol* **439**:46-64.
- Ciacchi C, Lind SE and Podolsky DK (1993) Transforming growth factor beta regulation of migration in wounded rat intestinal epithelial monolayers. *Gastroenterology* **105**:93-101.
- Ciruela F, Blanco J, Canelo EI, Lluis C, Franco R and Mallol J (1994) Solubilization and molecular characterization of the nitrobenzylthioinosine binding sites from pig kidney brush-border membranes. *Biochim Biophys Acta* **1191**:94-102.
- Clancy JP, Ruiz FE and Sorscher EJ (1999) Adenosine and its nucleotides activate wild-type and R117H CFTR through an A2B receptor-coupled pathway. *Am J Physiol* **276**:C361-9.
- Coe IR, Yao L, Diamond I and Gordon AS (1996) The role of protein kinase C in cellular tolerance to ethanol. *J Biol Chem* **271**:29468-72.
- Coe IR, Zhang Y, McKenzie T, Naydenova Z (2002) PKC regulation of human equilibrative nucleoside transporter, hENT1. *FEBS Lett* **517**: 201-5
- Collier AC, Coombs RW, Schoenfeld DA, Bassett RL, Timpone J, Baruch A, Jones M, Facey K, Whitacre C, McAuliffe VJ, Friedman HM, Merigan TC, Reichman RC, Hooper C and Corey L (1996) Treatment of human immunodeficiency virus infection with saquinavir, zidovudine, and zalcitabine. AIDS Clinical Trials Group. *N Engl J Med* **334**:1011-7.
- Corpe CP, Bovelander FJ, Hoekstra JH and Burant CF (1998) The small intestinal fructose transporters: site of dietary perception and evidence for diurnal and fructose sensitive control elements. *Biochim Biophys Acta* **1402**:229-38.
- Craig JE, Zhang Y and Gallagher MP (1994) Cloning of the nupC gene of Escherichia coli encoding a nucleoside transport system, and identification of an adjacent insertion element, IS 186. *Mol Microbiol* **11**:1159-68.
- Crawford CR, Patel DH, Naeve C and Belt JA (1998) Cloning of the human equilibrative, nitrobenzylmercaptopurine riboside (NBMPR)-insensitive nucleoside transporter ei by functional expression in a transport-deficient cell line. *J Biol Chem* **273**:5288-93.
- Crute BE, Seefeld K, Gamble J, Kemp BE and Witters LA (1998) Functional domains of the alpha1 catalytic subunit of the AMP-activated protein kinase. *J Biol Chem* **273**:35347-54.

D

- Dagnino L and Paterson AR (1990) Sodium-dependent and equilibrative nucleoside transport systems in L1210 mouse leukemia cells: effect of inhibitors of equilibrative systems on the content and retention of nucleosides. *Cancer Res* **50**:6549-53.
- Daniel T and Carling D (2002) Functional analysis of mutations in the gamma 2 subunit of AMP-activated protein kinase associated with cardiac hypertrophy and Wolff-Parkinson-White syndrome. *J Biol Chem* **277**:51017-24.
- Davies SP, Hawley SA, Woods A, Carling D, Haystead TA and Hardie DG (1994) Purification of the AMP-activated protein kinase on ATP-gamma-sepharose and analysis of its subunit structure. *Eur J Biochem* **223**:351-7.
- Davies SP, Helps NR, Cohen PT and Hardie DG (1995) 5'-AMP inhibits dephosphorylation, as well as promoting phosphorylation, of the AMP-activated protein kinase. Studies using bacterially expressed human protein phosphatase-2C alpha and native bovine protein phosphatase-2AC. *FEBS Lett* **377**:421-5.
- del Santo B, Tarafa G, Felipe A, Casado FJ and Pastor-Anglada M (2001) Developmental regulation of the concentrative nucleoside transporters CNT1 and CNT2 in rat liver. *J Hepatol* **34**:873-80.
- del Santo B, Valdes R, Mata J, Felipe A, Casado FJ and Pastor-Anglada M (1998) Differential expression and regulation of nucleoside transport systems in rat liver parenchymal and hepatoma cells. *Hepatology* **28**:1504-11.

- Dignass AU and Podolsky DK (1993) Cytokine modulation of intestinal epithelial cell restitution: central role of transforming growth factor beta. *Gastroenterology* **105**:1323-32.
- Dignass AU, Stow JL and Babyatsky MW (1996) Acute epithelial injury in the rat small intestine in vivo is associated with expanded expression of transforming growth factor alpha and beta. *Gut* **38**:687-93.
- Dionne S, D'Agata ID, Ruemmele FM, Levy E, St-Louis J, Srivastava AK, Levesque D and Seidman EG (1998) Tyrosine kinase and MAPK inhibition of TNF-alpha- and EGF-stimulated IEC-6 cell growth. *Biochem Biophys Res Commun* **242**:146-50.
- Doell RG, Rosen G and Kretchmer N (1965) Immunochemical studies of intestinal disaccharidases during normal and precocious development. *Proc Natl Acad Sci U S A* **54**:1268-73.
- Doherty AJ and Jarvis SM (1993) Na⁺-dependent and -independent uridine uptake in an established renal epithelial cell line, OK, from the opossum kidney. *Biochim Biophys Acta* **1147**:214-22.
- Dragan Y, Valdes R, Gomez-Angelats M, Felipe A, Javier Casado F, Pitot H and Pastor-Anglada M (2000) Selective loss of nucleoside carrier expression in rat hepatocarcinomas. *Hepatology* **32**:239-46.
- Duflot S, Calvo M, Casado FJ, Enrich C and Pastor-Anglada M (2002) Concentrative nucleoside transporter (rCNT1) is targeted to the apical membrane through the hepatic transcytotic pathway. *Exp Cell Res* **281**:77-85.
- Duflot S, Riera B, Fernandez-Veledo S, Casado V, Norman RI, Casado FJ, Lluis C, Franco R and Pastor-Anglada M (2004) ATP-sensitive K⁺ channels regulate the concentrative adenosine transporter CNT2 following activation by A(1) adenosine receptors. *Mol Cell Biol* **24**:2710-9.

E

- Endres CJ and Unadkat JD (2005) Residues Met89 and Ser160 in the human equilibrative nucleoside transporter 1 affect its affinity for adenosine, guanosine, S6-(4-nitrobenzyl)-mercaptopurine riboside, and dipyridamole. *Mol Pharmacol* **67**:837-44.
- Engel K, Zhou M and Wang J (2004) Identification and characterization of a novel monoamine transporter in the human brain. *J Biol Chem* **279**:50042-9.
- Erickson RH, Gum JR, Jr., Lindstrom MM, McKean D and Kim YS (1995) Regional expression and dietary regulation of rat small intestinal peptide and amino acid transporter mRNAs. *Biochem Biophys Res Commun* **216**:249-57.
- Estruch F, Treitel MA, Yang X and Carlson M (1992) N-terminal mutations modulate yeast SNF1 protein kinase function. *Genetics* **132**:639-50.

F

- Farre X, Guillen-Gomez E, Sanchez L, Hardisson D, Plaza Y, Lloberas J, Casado FJ, Palacios J and Pastor-Anglada M (2004) Expression of the nucleoside-derived drug transporters hCNT1, hENT1 and hENT2 in gynecologic tumors. *Int J Cancer* **112**:959-66.
- Feil W, Wenzl E, Vattay P, Starlinger M, Sogukoglu T and Schiessel R (1987) Repair of rabbit duodenal mucosa after acid injury in vivo and in vitro. *Gastroenterology* **92**:1973-86.
- Felipe A, Ferrer-Martinez A, Casado FJ and Pastor-Anglada M (1997) Expression of sodium-dependent purine nucleoside carrier (SPNT) mRNA correlates with nucleoside transport activity in rat liver. *Biochem Biophys Res Commun* **233**:572-5.
- Felipe A, Valdes R, Santo B, Lloberas J, Casado J and Pastor-Anglada M (1998) Na⁺-dependent nucleoside transport in liver: two different isoforms from the same gene family are expressed in liver cells. *Biochem J* **330 (Pt 2)**:997-1001.
- Fernandez-Veledo S, Valdes R, Wallenius V, Casado FJ and Pastor-Anglada M (2004) Up-regulation of the high-affinity pyrimidine-preferring nucleoside transporter concentrative nucleoside transporter 1 by tumor necrosis factor-alpha and interleukin-6 in liver parenchymal cells. *J Hepatol* **41**:538-44.
- Ferraris RP, Villenas SA, Hirayama BA and Diamond J (1992) Effect of diet on glucose transporter site density along the intestinal crypt-villus axis. *Am J Physiol* **262**:G1060-8.
- Ferre P, Azzout-Marniche D and Foufelle F (2003) AMP-activated protein kinase and hepatic genes involved in glucose metabolism. *Biochem Soc Trans* **31**:220-3.

- Ferruzza S, Ranaldi G, Di Girolamo M and Sambuy Y (1995) The transport of lysine across monolayers of human cultured intestinal cells (Caco-2) depends on Na(+) -dependent and Na(+) -independent mechanisms on different plasma membrane domains. *J Nutr* **125**:2577-85.
- Fischer B, Yefidoff R, Major DT, Rutman-Halili I, Shneyvays V, Zinman T, Jacobson KA and Shainberg A (1999) Characterization of "mini-nucleotides" as P2X receptor agonists in rat cardiomyocyte cultures. An integrated synthetic, biochemical, and theoretical study. *J Med Chem* **42**:2685-96.
- Flanagan SA, Gandhi V, Secrist JA, 3rd and Meckling KA (2003) The novel nucleoside transport system exhibited by NB4 cells, csg, transports deoxyguanosine analogues, including ara-G. *Biochem Pharmacol* **66**:733-7.
- Flanagan SA and Meckling-Gill KA (1997) Characterization of a novel Na⁺-dependent, guanosine-specific, nitrobenzylthioinosine-sensitive transporter in acute promyelocytic leukemia cells. *J Biol Chem* **272**:18026-32.
- Fontenelle LJ and Henderson JF (1969) An enzymatic basis for the inability of erythrocytes to synthesize purine ribonucleotides de novo. *Biochim Biophys Acta* **177**:175-6.
- Foretz M, Carling D, Guichard C, Ferre P and Foufelle F (1998) AMP-activated protein kinase inhibits the glucose-activated expression of fatty acid synthase gene in rat hepatocytes. *J Biol Chem* **273**:14767-71.
- Franco R, Centelles JJ and Kinne RK (1990) Further characterization of adenosine transport in renal brush-border membranes. *Biochim Biophys Acta* **1024**:241-8.
- Freathy C, Brown DG, Roberts RA and Cain K (2000) Transforming growth factor-beta(1) induces apoptosis in rat FaO hepatoma cells via cytochrome c release and oligomerization of Apaf-1 to form a approximately 700-kd apoptosome caspase-processing complex. *Hepatology* **32**:750-60.
- Fryer LG, Parbu-Patel A and Carling D (2002) The Anti-diabetic drugs rosiglitazone and metformin stimulate AMP-activated protein kinase through distinct signaling pathways. *J Biol Chem* **277**:25226-32.

G

- Galmarini CM, Mackey JR and Dumontet C (2001) Nucleoside analogues: mechanisms of drug resistance and reversal strategies. *Leukemia* **15**:875-90.
- Galmarini CM, Thomas X, Calvo F, Rousselot P, Jafaari AE, Cros E and Dumontet C (2002a) Potential mechanisms of resistance to cytarabine in AML patients. *Leuk Res* **26**:621-9.
- Galmarini CM, Thomas X, Calvo F, Rousselot P, Rabilloud M, El Jaffari A, Cros E and Dumontet C (2002b) In vivo mechanisms of resistance to cytarabine in acute myeloid leukaemia. *Br J Haematol* **117**:860-8.
- Gandhi V, Plunkett W, Rodriguez CO, Jr., Nowak BJ, Du M, Ayres M, Kisor DF, Mitchell BS, Kurtzberg J and Keating MJ (1998) Compound GW506U78 in refractory hematologic malignancies: relationship between cellular pharmacokinetics and clinical response. *J Clin Oncol* **16**:3607-15.
- Gasser T, Moyer JD and Handschumacher RE (1981) Novel single-pass exchange of circulating uridine in rat liver. *Science* **213**:777-8.
- Gati WP, Paterson AR, Belch AR, Chlumecky V, Larratt LM, Mant MJ and Turner AR (1998) Es nucleoside transporter content of acute leukemia cells: role in cell sensitivity to cytarabine (araC). *Leuk Lymphoma* **32**:45-54.
- Gati WP, Paterson AR, Larratt LM, Turner AR and Belch AR (1997) Sensitivity of acute leukemia cells to cytarabine is a correlate of cellular es nucleoside transporter site content measured by flow cytometry with SAENTA-fluorescein. *Blood* **90**:346-53.
- Giacomini KM, Markovich D, Werner A, Biber J, Wu X and Murer H (1994) Expression of a renal Na(+) -nucleoside cotransport system (N2) in Xenopus laevis oocytes. *Pflugers Arch* **427**:381-3.
- Gianneccini M, Matteucci M, Pesi R, Sgarrella F, Tozzi MG and Camici M (2005) Uptake and utilization of nucleosides for energy repletion. *Int J Biochem Cell Biol* **37**:797-808.
- Goke M, Kanai M, Lynch-Devaney K and Podolsky DK (1998) Rapid mitogen-activated protein kinase activation by transforming growth factor alpha in wounded rat intestinal epithelial cells. *Gastroenterology* **114**:697-705.

- Golin-Bisello F, Bradbury N and Ameen N (2005) STa and cGMP stimulate CFTR translocation to the surface of villus enterocytes in rat jejunum and is regulated by protein kinase G. *Am J Physiol Cell Physiol* **289**:C708-16.
- Gomez-Angelats M, Ruiz-Montasell B, Felipe A, Marin JJ, Casado FJ and Pastor-Anglada M (1995) Effect of protein malnutrition on neutral amino acid transport by rat hepatocytes during development. *Am J Physiol* **268**:E368-74.
- Goodyear LJ and Kahn BB (1998) Exercise, glucose transport, and insulin sensitivity. *Annu Rev Med* **49**:235-61.
- Graves LM, Guy HI, Kozlowski P, Huang M, Lazarowski E, Pope RM, Collins MA, Dahlstrand EN, Earp HS, 3rd and Evans DR (2000) Regulation of carbamoyl phosphate synthetase by MAP kinase. *Nature* **403**:328-32.
- Gray JH, Owen RP and Giacomini KM (2004) The concentrative nucleoside transporter family, SLC28. *Pflugers Arch* **447**:728-34.
- Greenberg N, Schumm DE and Webb TE (1977) Uridine kinase activities and pyrimidine nucleoside phosphorylation in fluoropyrimidine-sensitive and -resistant cell lines of the Novikoff hepatoma. *Biochem J* **164**:379-87.
- Grem JL and Fischer PH (1989) Enhancement of 5-fluorouracil's anticancer activity by dipyridamole. *Pharmacol Ther* **40**:349-71.
- Griffith DA and Jarvis SM (1991) Expression of sodium-dependent nucleoside transporters in Xenopus oocytes. *Adv Exp Med Biol* **309A**:431-4.
- Griffith DA and Jarvis SM (1996) Nucleoside and nucleobase transport systems of mammalian cells. *Biochim Biophys Acta* **1286**:153-81.
- Griffiths M, Beaumont N, Yao SY, Sundaram M, Boumah CE, Davies A, Kwong FY, Coe I, Cass CE, Young JD and Baldwin SA (1997a) Cloning of a human nucleoside transporter implicated in the cellular uptake of adenosine and chemotherapeutic drugs. *Nat Med* **3**:89-93.
- Griffiths M, Yao SY, Abidi F, Phillips SE, Cass CE, Young JD and Baldwin SA (1997b) Molecular cloning and characterization of a nitrobenzylthioinosine-insensitive (ei) equilibrative nucleoside transporter from human placenta. *Biochem J* **328** (Pt 3):739-43.
- Gu JJ, Gathy K, Santiago L, Chen E, Huang M, Graves LM and Mitchell BS (2003) Induction of apoptosis in IL-3-dependent hematopoietic cell lines by guanine nucleotide depletion. *Blood* **101**:4958-65.
- Guillen-Gomez E, Calbet M, Casado J, de Lecea L, Soriano E, Pastor-Anglada M and Burgaya F (2004) Distribution of CNT2 and ENT1 transcripts in rat brain: selective decrease of CNT2 mRNA in the cerebral cortex of sleep-deprived rats. *J Neurochem* **90**:883-93.
- Gutierrez MM and Giacomini KM (1993) Substrate selectivity, potential sensitivity and stoichiometry of Na⁺-nucleoside transport in brush border membrane vesicles from human kidney. *Biochim Biophys Acta* **1149**:202-8.

H

- Hack SP and Christie MJ (2003) Adaptations in adenosine signaling in drug dependence: therapeutic implications. *Crit Rev Neurobiol* **15**:235-74.
- Haidari M, Leung N, Mahbub F, Uffelman KD, Kohen-Avramoglu R, Lewis GF and Adeli K (2002) Fasting and postprandial overproduction of intestinally derived lipoproteins in an animal model of insulin resistance. Evidence that chronic fructose feeding in the hamster is accompanied by enhanced intestinal de novo lipogenesis and ApoB48-containing lipoprotein overproduction. *J Biol Chem* **277**:31646-55.
- Hamilton SR, O'Donnell JB, Jr., Hammet A, Stapleton D, Habinowski SA, Means AR, Kemp BE and Witters LA (2002) AMP-activated protein kinase kinase: detection with recombinant AMPK alpha1 subunit. *Biochem Biophys Res Commun* **293**:892-8.
- Hamilton SR, Yao SY, Ingram JC, Hadden DA, Ritzel MW, Gallagher MP, Henderson PJ, Cass CE, Young JD and Baldwin SA (2001) Subcellular distribution and membrane topology of the mammalian concentrative Na⁺-nucleoside cotransporter rCNT1. *J Biol Chem* **276**:27981-8.
- Hammond JR, Lee S and Ferguson PJ (1999) [3H]gemcitabine uptake by nucleoside transporters in a human head and neck squamous carcinoma cell line. *J Pharmacol Exp Ther* **288**:1185-91.

- Hancock DL and Coupar IM (1995) Functional characterization of the adenosine receptor mediating inhibition of intestinal secretion. *Br J Pharmacol* **114**:152-6.
- Hanks SK, Quinn AM and Hunter T (1988) The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* **241**:42-52.
- Hanna NN, Mauceri HJ, Wayne JD, Hallahan DE, Kufe DW and Weichselbaum RR (1997) Virally directed cytosine deaminase/5-fluorocytosine gene therapy enhances radiation response in human cancer xenografts. *Cancer Res* **57**:4205-9.
- Hansen RW, Ranelli PL and Ried LD (1995) Stigma, conflict, and the approval of AIDS drugs. *J Drug Issues* **25**:129-39.
- Hardie DG (1999) Roles of the AMP-activated/SNF1 protein kinase family in the response to cellular stress. *Biochem Soc Symp* **64**:13-27.
- Hardie DG (2003) Minireview: the AMP-activated protein kinase cascade: the key sensor of cellular energy status. *Endocrinology* **144**:5179-83.
- Hardie DG (2004) The AMP-activated protein kinase pathway--new players upstream and downstream. *J Cell Sci* **117**:5479-87.
- Hardie DG and Carling D (1997) The AMP-activated protein kinase--fuel gauge of the mammalian cell? *Eur J Biochem* **246**:259-73.
- Hardie DG and Hawley SA (2001) AMP-activated protein kinase: the energy charge hypothesis revisited. *Bioessays* **23**:1112-9.
- Hardin JA, Wong JK, Cheeseman CI and Gall DG (1996) Effect of luminal epidermal growth factor on enterocyte glucose and proline transport. *Am J Physiol* **271**:G509-15.
- Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Makela TP, Alessi DR and Hardie DG (2003) Complexes between the LKB1 tumor suppressor, STRADalpha/beta and MO25alpha/beta are upstream kinases in the AMP-activated protein kinase cascade. *J Biol* **2**:28.
- Hawley SA, Davison M, Woods A, Davies SP, Beri RK, Carling D and Hardie DG (1996) Characterization of the AMP-activated protein kinase kinase from rat liver and identification of threonine 172 as the major site at which it phosphorylates AMP-activated protein kinase. *J Biol Chem* **271**:27879-87.
- Hawley SA, Selbert MA, Goldstein EG, Edelman AM, Carling D and Hardie DG (1995) 5'-AMP activates the AMP-activated protein kinase cascade, and Ca²⁺/calmodulin activates the calmodulin-dependent protein kinase I cascade, via three independent mechanisms. *J Biol Chem* **270**:27186-91.
- Hayashi T, Hirshman MF, Kurth EJ, Winder WW and Goodyear LJ (1998) Evidence for 5' AMP-activated protein kinase mediation of the effect of muscle contraction on glucose transport. *Diabetes* **47**:1369-73.
- He Y, Chu SH and Walker WA (1993) Nucleotide supplements alter proliferation and differentiation of cultured human (Caco-2) and rat (IEC-6) intestinal epithelial cells. *J Nutr* **123**:1017-27.
- He Y, Sanderson IR and Walker WA (1994) Uptake, transport and metabolism of exogenous nucleosides in intestinal epithelial cell cultures. *J Nutr* **124**:1942-9.
- Hemminki A, Markie D, Tomlinson I, Avizienyte E, Roth S, Loukola A, Bignell G, Warren W, Aminoff M, Hoglund P, Jarvinen H, Kristo P, Pelin K, Ridanpaa M, Salovaara R, Toro T, Bodmer W, Olschwang S, Olsen AS, Stratton MR, de la Chapelle A and Aaltonen LA (1998) A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* **391**:184-7.
- Hendriks G, Koudijs M, van Balkom BW, Oorschot V, Klumperman J, Deen PM and van der Sluijs P (2004) Glycosylation is important for cell surface expression of the water channel aquaporin-2 but is not essential for tetramerization in the endoplasmic reticulum. *J Biol Chem* **279**:2975-83.
- Herzfeld A and Raper SM (1979) Uridine kinase activities in developing, adult and neoplastic rat tissues. *Biochem J* **182**:771-8.
- Hirano M, Iwakiri R, Fujimoto K, Sakata H, Ohyama T, Sakai T, Joh T and Itoh M (1995) Epidermal growth factor enhances repair of rat intestinal mucosa damaged by oral administration of methotrexate. *J Gastroenterol* **30**:169-76.
- Holstege A, Gengenbacher HM, Jehle L and Hoppmann J (1991) Facilitated diffusion and sodium-dependent transport of purine and pyrimidine nucleosides in rat liver. *Hepatology* **14**:373-80.

- Hong M, Schlichter L and Bendayan R (2001) A novel zidovudine uptake system in microglia. *J Pharmacol Exp Ther* **296**:141-9.
- Horman S, Browne G, Krause U, Patel J, Vertommen D, Bertrand L, Lavoinne A, Hue L, Proud C and Rider M (2002) Activation of AMP-activated protein kinase leads to the phosphorylation of elongation factor 2 and an inhibition of protein synthesis. *Curr Biol* **12**:1419-23.
- Huang P, Chubb S, Hertel LW, Grindey GB and Plunkett W (1991) Action of 2',2'-difluorodeoxycytidine on DNA synthesis. *Cancer Res* **51**:6110-7.
- Huang P, Robertson LE, Wright S and Plunkett W (1995) High molecular weight DNA fragmentation: a critical event in nucleoside analogue-induced apoptosis in leukemia cells. *Clin Cancer Res* **1**:1005-13.
- Huang QQ, Harvey CM, Paterson AR, Cass CE and Young JD (1993) Functional expression of Na(+) -dependent nucleoside transport systems of rat intestine in isolated oocytes of *Xenopus laevis*. Demonstration that rat jejunum expresses the purine-selective system N1 (cif) and a second, novel system N3 having broad specificity for purine and pyrimidine nucleosides. *J Biol Chem* **268**:20613-9.
- Huang QQ, Yao SY, Ritzel MW, Paterson AR, Cass CE and Young JD (1994) Cloning and functional expression of a complementary DNA encoding a mammalian nucleoside transport protein. *J Biol Chem* **269**:17757-60.
- Hubert A, Husson A, Chedeville A and Lavoinne A (2000) AMP-activated protein kinase counteracted the inhibitory effect of glucose on the phosphoenolpyruvate carboxykinase gene expression in rat hepatocytes. *FEBS Lett* **481**:209-12.
- Hutber CA, Hardie DG and Winder WW (1997) Electrical stimulation inactivates muscle acetyl-CoA carboxylase and increases AMP-activated protein kinase. *Am J Physiol* **272**:E262-6.
- Hyde RJ, Cass CE, Young JD and Baldwin SA (2001) The ENT family of eukaryote nucleoside and nucleobase transporters: recent advances in the investigation of structure/function relationships and the identification of novel isoforms. *Mol Membr Biol* **18**:53-63.

I

- Iannoli P, Miller JH, Ryan CK and Sax HC (1998a) Glucocorticoids upregulate intestinal nutrient transport in a time-dependent and substrate-specific fashion. *J Gastrointest Surg* **2**:449-57.
- Iannoli P, Miller JH and Sax HC (1998b) Epidermal growth factor and human growth hormone induce two sodium-dependent arginine transport systems after massive enterectomy. *JPEN J Parenter Enteral Nutr* **22**:326-30.
- Imamura K, Ogura T, Kishimoto A, Kaminishi M and Esumi H (2001) Cell cycle regulation via p53 phosphorylation by a 5'-AMP activated protein kinase activator, 5-aminoimidazole-4-carboxamide-1-beta-D-ribofuranoside, in a human hepatocellular carcinoma cell line. *Biochem Biophys Res Commun* **287**:562-7.
- Inoki K, Zhu T and Guan KL (2003) TSC2 mediates cellular energy response to control cell growth and survival. *Cell* **115**:577-90.
- Inui K, Quaroni A, Tillotson LG and Isselbacher KJ (1980) Amino acid and hexose transport by cultured crypt cells from rat small intestine. *Am J Physiol* **239**:C190-6.
- Iseki K, Sugawara M, Fujiwara T, Naasani I, Kobayashi M and Miyazaki K (1996) Transport mechanisms of nucleosides and the derivative, 6-mercaptopurine riboside across rat intestinal brush-border membranes. *Biochim Biophys Acta* **1278**:105-10.
- Isselbacher KJ (1974) The intestinal cell surface: some properties of normal, undifferentiated, and malignant cells. *Ann Intern Med* **81**:681-6.
- Iwasa Y, Iwasa M, Ohmori Y, Fukutomi T and Ogoshi S (2000) The effect of the administration of nucleosides and nucleotides for parenteral use. *Nutrition* **16**:598-602.
- Iwasaki H, Huang P, Keating MJ and Plunkett W (1997) Differential incorporation of ara-C, gemcitabine, and fludarabine into replicating and repairing DNA in proliferating human leukemia cells. *Blood* **90**:270-8.

J

- Jakobs ES, Van Os-Corby DJ and Paterson AR (1990) Expression of sodium-linked nucleoside transport activity in monolayer cultures of IEC-6 intestinal epithelial cells. *J Biol Chem* **265**:22210-6.
- Jakobsen SN, Hardie DG, Morrice N and Tornqvist HE (2001) 5'-AMP-activated protein kinase phosphorylates IRS-1 on Ser-789 in mouse C2C12 myotubes in response to 5-aminoimidazole-4-carboxamide riboside. *J Biol Chem* **276**:46912-6.
- Jakobsson B, el Hag IA, Erichsen C, Christensson PI, Jonsson PE and Stenram U (1989) Modulation of 5-fluorouracil and 5-fluorouridine toxicity by membrane transport inhibitors in normal tissues of rats with liver adenocarcinoma. *Anticancer Res* **9**:285-90.
- Jamieson GP, Snook MB, Bradley TR, Bertoncello I and Wiley JS (1989) Transport and metabolism of 1-beta-D-arabinofuranosylcytosine in human ovarian adenocarcinoma cells. *Cancer Res* **49**:309-13.
- Jarvis MF, Yu H, McGaraughty S, Wismer CT, Mikusa J, Zhu C, Chu K, Kohlhaas K, Cowart M, Lee CH, Stewart AO, Cox BF, Polakowski J and Kowaluk EA (2002) Analgesic and anti-inflammatory effects of A-286501, a novel orally active adenosine kinase inhibitor. *Pain* **96**:107-18.
- Jarvis SM and Griffith DA (1991) Expression of the rabbit intestinal N₂ Na⁺/nucleoside transporter in *Xenopus laevis* oocytes. *Biochem J* **278 (Pt 2)**:605-7.
- Jarvis SM and Young JD (1981) Extraction and partial purification of the nucleoside-transport system from human erythrocytes based on the assay of nitrobenzylthioinosine-binding activity. *Biochem J* **194**:331-9.
- Jarvis SM, Young JD and Ellory JC (1980) Nucleoside transport in human erythrocytes. Apparent molecular weight of the nitrobenzylthioinosine-binding complex estimated by radiation-inactivation analysis. *Biochem J* **190**:373-6.
- Jenne DE, Reimann H, Nezu J, Friedel W, Loff S, Jeschke R, Muller O, Back W and Zimmer M (1998) Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat Genet* **18**:38-43.
- Jimenez AI, Fernandez P, Dominguez O, Dopazo A and Sanchez-Cespedes M (2003) Growth and molecular profile of lung cancer cells expressing ectopic LKB1: down-regulation of the phosphatidylinositol 3'-phosphate kinase/PTEN pathway. *Cancer Res* **63**:1382-8.

K

- Kanai Y, Segawa H, Chairoungdua A, Kim JY, Kim DK, Matsuo H, Cha SH and Endou H (2000) Amino acid transporters: molecular structure and physiological roles. *Nephrol Dial Transplant* **15 Suppl 6**:9-10.
- Kashlan OB and Cooperman BS (2003) Comprehensive model for allosteric regulation of mammalian ribonucleotide reductase: refinements and consequences. *Biochemistry* **42**:1696-706.
- Katsura T and Inui K (2003) Intestinal absorption of drugs mediated by drug transporters: mechanisms and regulation. *Drug Metab Pharmacokinet* **18**:1-15.
- Kawaguchi T, Osatomi K, Yamashita H, Kabashima T and Uyeda K (2002) Mechanism for fatty acid "sparing" effect on glucose-induced transcription: regulation of carbohydrate-responsive element-binding protein by AMP-activated protein kinase. *J Biol Chem* **277**:3829-35.
- Kishi K, Tanaka T, Igawa M, Takase S and Goda T (1999) Sucrase-isomaltase and hexose transporter gene expressions are coordinately enhanced by dietary fructose in rat jejunum. *J Nutr* **129**:953-6.
- Kiss A, Farah K, Kim J, Garriock RJ, Drysdale TA and Hammond JR (2000) Molecular cloning and functional characterization of inhibitor-sensitive (mENT1) and inhibitor-resistant (mENT2) equilibrative nucleoside transporters from mouse brain. *Biochem J* **352 Pt 2**:363-72.
- Kolassa N, Stengg R and Turnheim K (1977) Salvage of adenosine, inosine, hypoxanthine, and adenine by the isolated epithelium of guinea pig jejunum. *Can J Physiol Pharmacol* **55**:1039-44.
- Kowaluk EA, Bhagwat SS and Jarvis MF (1998) Adenosine kinase inhibitors. *Curr Pharm Des* **4**:403-16.

- Koyama SY and Podolsky DK (1989) Differential expression of transforming growth factors alpha and beta in rat intestinal epithelial cells. *J Clin Invest* **83**:1768-73.
- Kroll K, Decking UK, Dreikorn K and Schrader J (1993) Rapid turnover of the AMP-adenosine metabolic cycle in the guinea pig heart. *Circ Res* **73**:846-56.
- Kurokawa M, Lynch K and Podolsky DK (1987) Effects of growth factors on an intestinal epithelial cell line: transforming growth factor beta inhibits proliferation and stimulates differentiation. *Biochem Biophys Res Commun* **142**:775-82.
- Kwong FY, Tse CM, Jarvis SM, Choy MY and Young JD (1987) Purification and reconstitution studies of the nucleoside transporter from pig erythrocytes. *Biochim Biophys Acta* **904**:105-16.

L

- Lacy ER (1988) Epithelial restitution in the gastrointestinal tract. *J Clin Gastroenterol* **10 Suppl 1**:S72-7.
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680-5.
- Lai Y, Bakken AH and Unadkat JD (2002) Simultaneous expression of hCNT1-CFP and hENT1-YFP in Madin-Darby canine kidney cells. Localization and vectorial transport studies. *J Biol Chem* **277**:37711-7.
- Lai Y, Lee EW, Ton CC, Vijay S, Zhang H and Unadkat JD (2005) Conserved residues F316 and G476 in the concentrative nucleoside transporter 1 (hCNT1) affect guanosine sensitivity and membrane expression, respectively. *Am J Physiol Cell Physiol* **288**:C39-45.
- Lai Y, Tse CM and Unadkat JD (2004) Mitochondrial expression of the human equilibrative nucleoside transporter 1 (hENT1) results in enhanced mitochondrial toxicity of antiviral drugs. *J Biol Chem* **279**:4490-7.
- Lajtha LG and Vane JR (1958) Dependence of bone marrow cells on the liver for purine supply. *Nature* **182**:191-2.
- Lang TT, Selner M, Young JD and Cass CE (2001) Acquisition of human concentrative nucleoside transporter 2 (hcnt2) activity by gene transfer confers sensitivity to fluoropyrimidine nucleosides in drug-resistant leukemia cells. *Mol Pharmacol* **60**:1143-52.
- Larrayoz IM, Casado FJ, Pastor-Anglada M and Lostao MP (2004) Electrophysiological characterization of the human Na⁺/nucleoside cotransporter 1 (hCNT1) and role of adenosine on hCNT1 function. *J Biol Chem* **279**:8999-9007.
- Lasley RD, Hegge JO, Noble MA and Mentzer RM, Jr. (1998) Comparison of interstitial fluid and coronary venous adenosine levels in vivo porcine myocardium. *J Mol Cell Cardiol* **30**:1137-47.
- Leclerc I, Kahn A and Doiron B (1998) The 5'-AMP-activated protein kinase inhibits the transcriptional stimulation by glucose in liver cells, acting through the glucose response complex. *FEBS Lett* **431**:180-4.
- Leclerc I, Lenzner C, Gourdon L, Vaulont S, Kahn A and Viollet B (2001) Hepatocyte nuclear factor-4alpha involved in type 1 maturity-onset diabetes of the young is a novel target of AMP-activated protein kinase. *Diabetes* **50**:1515-21.
- Lee CW, Cheeseman CI and Jarvis SM (1988) Na⁺- and K⁺-dependent uridine transport in rat renal brush-border membrane vesicles. *Biochim Biophys Acta* **942**:139-49.
- Lee K, Li B, Xi X, Suh Y and Martin RJ (2005) Role of neuronal energy status in the regulation of adenosine 5'-monophosphate-activated protein kinase, orexigenic neuropeptides expression, and feeding behavior. *Endocrinology* **146**:3-10.
- Leibovitz A (1963) The Growth and Maintenance of Tissue-Cell Cultures in Free Gas Exchange with the Atmosphere. *Am J Hyg* **78**:173-80.
- Lizcano JM, Goransson O, Toth R, Deak M, Morrice NA, Boudeau J, Hawley SA, Udd L, Makela TP, Hardie DG and Alessi DR (2004) LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. *Embo J* **23**:833-43.
- Lloyd HG, Spence I and Johnston GA (1988) Protective effect of adenosine against hypoxic damage. Synaptic depression and adenosine. *Trends Pharmacol Sci* **9**:352.

- Lochhead PA, Salt IP, Walker KS, Hardie DG and Sutherland C (2000) 5-aminoimidazole-4-carboxamide riboside mimics the effects of insulin on the expression of the 2 key gluconeogenic genes PEPCK and glucose-6-phosphatase. *Diabetes* **49**:896-903.
- Loewen SK, Ng AM, Yao SY, Cass CE, Baldwin SA and Young JD (1999) Identification of amino acid residues responsible for the pyrimidine and purine nucleoside specificities of human concentrative Na(+) nucleoside cotransporters hCNT1 and hCNT2. *J Biol Chem* **274**:24475-84.
- Lostao MP, Mata JF, Larrayoz IM, Inzillo SM, Casado FJ and Pastor-Anglada M (2000) Electrogenic uptake of nucleosides and nucleoside-derived drugs by the human nucleoside transporter 1 (hCNT1) expressed in *Xenopus laevis* oocytes. *FEBS Lett* **481**:137-40.
- Lund P, Cornell NW and Krebs HA (1975) Effect of adenosine on the adenine nucleotide content and metabolism of hepatocytes. *Biochem J* **152**:593-9.
- Luo Z, Saha AK, Xiang X and Ruderman NB (2005) AMPK, the metabolic syndrome and cancer. *Trends Pharmacol Sci* **26**:69-76.

M

- Mackey JR, Galmarini CM, Graham KA, Joy AA, Delmer A, Dabbagh L, Glubrecht D, Jewell LD, Lai R, Lang T, Hanson J, Young JD, Merle-Beral H, Binet JL, Cass CE and Dumontet C (2005) Quantitative analysis of nucleoside transporter and metabolism gene expression in chronic lymphocytic leukemia (CLL): identification of fludarabine-sensitive and -insensitive populations. *Blood* **105**:767-74.
- Mackey JR, Jennings LL, Clarke ML, Santos CL, Dabbagh L, Vsianska M, Koski SL, Coupland RW, Baldwin SA, Young JD and Cass CE (2002) Immunohistochemical variation of human equilibrative nucleoside transporter 1 protein in primary breast cancers. *Clin Cancer Res* **8**:110-6.
- Mackey JR, Mani RS, Selner M, Mowles D, Young JD, Belt JA, Crawford CR and Cass CE (1998) Functional nucleoside transporters are required for gemcitabine influx and manifestation of toxicity in cancer cell lines. *Cancer Res* **58**:4349-57.
- Mackey JR, Yao SY, Smith KM, Karpinski E, Baldwin SA, Cass CE and Young JD (1999) Gemcitabine transport in *xenopus* oocytes expressing recombinant plasma membrane mammalian nucleoside transporters. *J Natl Cancer Inst* **91**:1876-81.
- Mangravite LM, Badagnani I and Giacomini KM (2003a) Nucleoside transporters in the disposition and targeting of nucleoside analogs in the kidney. *Eur J Pharmacol* **479**:269-81.
- Mangravite LM and Giacomini KM (2003) Sorting of rat SPNT in renal epithelium is independent of N-glycosylation. *Pharm Res* **20**:319-23.
- Mangravite LM, Lipschutz JH, Mostov KE and Giacomini KM (2001) Localization of GFP-tagged concentrative nucleoside transporters in a renal polarized epithelial cell line. *Am J Physiol Renal Physiol* **280**:F879-85.
- Mangravite LM, Xiao G and Giacomini KM (2003b) Localization of human equilibrative nucleoside transporters, hENT1 and hENT2, in renal epithelial cells. *Am J Physiol Renal Physiol* **284**:F902-10.
- Marmorstein AD, Csaky KG, Baffi J, Lam L, Rahaal F and Rodriguez-Boulan E (2000) Saturation of, and competition for entry into, the apical secretory pathway. *Proc Natl Acad Sci U S A* **97**:3248-53.
- Mata JF, Garcia-Manteiga JM, Lostao MP, Fernandez-Veledo S, Guillen-Gomez E, Larrayoz IM, Lloberas J, Casado FJ and Pastor-Anglada M (2001) Role of the human concentrative nucleoside transporter (hCNT1) in the cytotoxic action of 5[Prime]-deoxy-5-fluorouridine, an active intermediate metabolite of capecitabine, a novel oral anticancer drug. *Mol Pharmacol* **59**:1542-8.
- Matsumoto K, Takao Y, Akazawa S, Yano M, Uotani S, Kawasaki E, Takino H, Yamasaki H, Okuno S, Yamaguchi Y and et al. (1993) Developmental change of facilitative glucose transporter expression in rat embryonal and fetal intestine. *Biochem Biophys Res Commun* **193**:1275-82.
- McCabe JB and Berthiaume LG (1999) Functional roles for fatty acylated amino-terminal domains in subcellular localization. *Mol Biol Cell* **10**:3771-86.

- McCormack SA, Viar MJ and Johnson LR (1992) Migration of IEC-6 cells: a model for mucosal healing. *Am J Physiol* **263**:G426-35.
- McCrimmon RJ, Fan X, Ding Y, Zhu W, Jacob RJ and Sherwin RS (2004) Potential role for AMP-activated protein kinase in hypoglycemia sensing in the ventromedial hypothalamus. *Diabetes* **53**:1953-8.
- McGinn CJ, Miller EM, Lindstrom MJ, Kunugi KA, Johnston PG and Kinsella TJ (1994) The role of cell cycle redistribution in radiosensitization: implications regarding the mechanism of fluorodeoxyuridine radiosensitization. *Int J Radiat Oncol Biol Phys* **30**:851-9.
- McGivan JD and Pastor-Anglada M (1994) Regulatory and molecular aspects of mammalian amino acid transport. *Biochem J* **299** (Pt 2):321-34.
- Meckling-Gill KA, Guilbert L and Cass CE (1993) CSF-1 stimulates nucleoside transport in S1 macrophages. *J Cell Physiol* **155**:530-8.
- Meisel H, Gunther S, Martin D and Schlimme E (1998) Apoptosis induced by modified ribonucleosides in human cell culture systems. *FEBS Lett* **433**:265-8.
- Mercader J, Gomez-Angelats M, del Santo B, Casado FJ, Felipe A and Pastor-Anglada M (1996) Nucleoside uptake in rat liver parenchymal cells. *Biochem J* **317** (Pt 3):835-42.
- Merriman RL, Hertel LW, Schultz RM, Houghton PJ, Houghton JA, Rutherford PG, Tanzer LR, Boder GB and Grindey GB (1996) Comparison of the antitumor activity of gemcitabine and ara-C in a panel of human breast, colon, lung and pancreatic xenograft models. *Invest New Drugs* **14**:243-7.
- Mikkaichi T, Suzuki T, Tanemoto M, Ito S and Abe T (2004) The organic anion transporter (OATP) family. *Drug Metab Pharmacokinet* **19**:171-9.
- Minokoshi Y, Alquier T, Furukawa N, Kim YB, Lee A, Xue B, Mu J, Foufelle F, Ferre P, Birnbaum MJ, Stuck BJ and Kahn BB (2004) AMP-kinase regulates food intake by responding to hormonal and nutrient signals in the hypothalamus. *Nature* **428**:569-74.
- Minokoshi Y, Kim YB, Peroni OD, Fryer LG, Muller C, Carling D and Kahn BB (2002) Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature* **415**:339-43.
- Miranda E, Romisch K and Lomas DA (2004) Mutants of neuroserpin that cause dementia accumulate as polymers within the endoplasmic reticulum. *J Biol Chem* **279**:28283-91.
- Miras-Portugal MT, Gualix J and Pintor J (1998) The neurotransmitter role of diadenosine polyphosphates. *FEBS Lett* **430**:78-82.
- Mitchellhill KI, Stapleton D, Gao G, House C, Michell B, Katsis F, Witters LA and Kemp BE (1994) Mammalian AMP-activated protein kinase shares structural and functional homology with the catalytic domain of yeast Snf1 protein kinase. *J Biol Chem* **269**:2361-4.
- Mohamedali KA, Guicherit OM, Kellems RE and Rudolph FB (1993) The highest levels of purine catabolic enzymes in mice are present in the proximal small intestine. *J Biol Chem* **268**:23728-33.
- Molina-Arcas M, Bellosillo B, Casado FJ, Montserrat E, Gil J, Colomer D and Pastor-Anglada M (2003) Fludarabine uptake mechanisms in B-cell chronic lymphocytic leukemia. *Blood* **101**:2328-34.
- Molina-Arcas M, Marce S, Villamor N, Huber-Ruano I, Casado FJ, Bellosillo B, Montserrat E, Gil J, Colomer D and Pastor-Anglada M (2005) Equilibrative nucleoside transporter-2 (hENT2) protein expression correlates with ex vivo sensitivity to fludarabine in chronic lymphocytic leukemia (CLL) cells. *Leukemia* **19**:64-8.
- Momparler RL, Eliopoulos N, Bovenzi V, Letourneau S, Greenbaum M and Cournoyer D (1996) Resistance to cytosine arabinoside by retrovirally mediated gene transfer of human cytidine deaminase into murine fibroblast and hematopoietic cells. *Cancer Gene Ther* **3**:331-8.
- Moreno F, Pastor-Anglada M, Hollenberg MD and Soley M (1989) Effects of epidermal growth factor (urogastrone) on gluconeogenesis, glucose oxidation, and glycogen synthesis in isolated rat hepatocytes. *Biochem Cell Biol* **67**:724-9.
- Mu J, Brozinick JT, Jr., Valladares O, Bucan M and Birnbaum MJ (2001) A role for AMP-activated protein kinase in contraction- and hypoxia-regulated glucose transport in skeletal muscle. *Mol Cell* **7**:1085-94.

Muoio DM, Seefeld K, Witters LA and Coleman RA (1999) AMP-activated kinase reciprocally regulates triacylglycerol synthesis and fatty acid oxidation in liver and muscle: evidence that sn-glycerol-3-phosphate acyltransferase is a novel target. *Biochem J* **338** (Pt 3):783-91.

N

- Ngo LY, Patil SD and Unadkat JD (2001) Ontogeny and longitudinal activity of Na(+)-nucleoside transporters in the human intestine. *Am J Physiol Gastrointest Liver Physiol* **280**:G475-81.
- Nicklin PL, Irwin WJ, Hassan IF, Mackay M and Dixon HB (1995) The transport of acidic amino acids and their analogues across monolayers of human intestinal absorptive (Caco-2) cells in vitro. *Biochim Biophys Acta* **1269**:176-86.
- Nielsen CU, Amstrup J, Steffansen B, Frokjaer S and Brodin B (2001) Epidermal growth factor inhibits glycylsarcosine transport and hPepT1 expression in a human intestinal cell line. *Am J Physiol Gastrointest Liver Physiol* **281**:G191-9.
- Novak DA, Carver JD and Barness LA (1994) Dietary nucleotides affect hepatic growth and composition in the weanling mouse. *JPEN J Parenter Enteral Nutr* **18**:62-6.

O

- Ohyanagi H, Nishimatsu S, Kanbara Y, Usami M and Saitoh Y (1989) Effects of nucleosides and a nucleotide on DNA and RNA syntheses by the salvage and de novo pathway in primary monolayer cultures of hepatocytes and hepatoma cells. *JPEN J Parenter Enteral Nutr* **13**:51-8.
- Oliver BL, Sha'afi RI and Hajjar JJ (1994) Transforming growth factor-alpha increases tyrosine phosphorylation of microtubule-associated protein kinase in a small intestinal crypt cell line (IEC-6). *Biochem J* **303** (Pt 2):455-60.
- Ortega MA, Nunez MC, Gil A and Sanchez-Pozo A (1995) Dietary nucleotides accelerate intestinal recovery after food deprivation in old rats. *J Nutr* **125**:1413-8.
- Osse N, Pearson JD, Yudilevich DL and Jarvis SM (1996) Hypoxanthine enters human vascular endothelial cells (ECV 304) via the nitrobenzylthioinosine-insensitive equilibrative nucleoside transporter. *Biochem J* **317** (Pt 3):843-8.

P

- Pagliassotti MJ and Cherrington AD (1992) Regulation of net hepatic glucose uptake in vivo. *Annu Rev Physiol* **54**:847-60.
- Pan M and Stevens BR (1995) Protein kinase C-dependent regulation of L-arginine transport activity in Caco-2 intestinal cells. *Biochim Biophys Acta* **1239**:27-32.
- Pang S, Urquhart P and Hooper NM (2004) N-glycans, not the GPI anchor, mediate the apical targeting of a naturally glycosylated, GPI-anchored protein in polarised epithelial cells. *J Cell Sci* **117**:5079-86.
- Park H, Kaushik VK, Constant S, Prentki M, Przybytkowski E, Ruderman NB and Saha AK (2002) Coordinate regulation of malonyl-CoA decarboxylase, sn-glycerol-3-phosphate acyltransferase, and acetyl-CoA carboxylase by AMP-activated protein kinase in rat tissues in response to exercise. *J Biol Chem* **277**:32571-7.
- Pasini FL, Guideri F, Ferber D, Galgano G, Bianchi A, Isidori S, De Giorgi L, Petri S, Capecchi PL and Di Perri T (1996) Pharmacological preconditioning of ischemic heart disease by low-dose dipyridamole. *Int J Cardiol* **56**:17-27.
- Pastor-Anglada M, Cano-Soldado P, Molina-Arcas M, Lostao MP, Larrayoz I, Martinez-Picado J and Casado FJ (2005) Cell entry and export of nucleoside analogues. *Virus Res* **107**:151-64.
- Pastor-Anglada M, Casado FJ, Valdes R, Mata J, Garcia-Manteiga J and Molina M (2001) Complex regulation of nucleoside transporter expression in epithelial and immune system cells. *Mol Membr Biol* **18**:81-5.
- Pastor-Anglada M, Felipe A and Casado FJ (1998a) Transport and mode of action of nucleoside derivatives used in chemical and antiviral therapies. *Trends Pharmacol Sci* **19**:424-30.

- Pastor-Anglada M, Felipe A, Casado FJ, del Santo B, Mata JF and Valdes R (1998b) Nucleoside transporters and liver cell growth. *Biochem Cell Biol* **76**:771-7.
- Pastor-Anglada M, Molina-Arcas M, Casado FJ, Bellosillo B, Colomer D and Gil J (2004) Nucleoside transporters in chronic lymphocytic leukaemia. *Leukemia* **18**:385-93.
- Patel DH, Crawford CR, Naeve CW and Belt JA (2000) Cloning, genomic organization and chromosomal localization of the gene encoding the murine sodium-dependent, purine-selective, concentrative nucleoside transporter (CNT2). *Gene* **242**:51-8.
- Patil SD and Unadkat JD (1997) Sodium-dependent nucleoside transport in the human intestinal brush-border membrane. *Am J Physiol* **272**:G1314-20.
- Patrick SM, Oakley GG, Dixon K and Turchi JJ (2005) DNA damage induced hyperphosphorylation of replication protein A. 2. Characterization of DNA binding activity, protein interactions, and activity in DNA replication and repair. *Biochemistry* **44**:8438-48.
- Pennycooke M, Chaudary N, Shuralyova I, Zhang Y and Coe IR (2001) Differential expression of human nucleoside transporters in normal and tumor tissue. *Biochem Biophys Res Commun* **280**:951-9.
- Petersen AJ, Brown RD, Gibson J, Pope B, Luo XF, Schutz L, Wiley JS and Joshua DE (1996) Nucleoside transporters, bcl-2 and apoptosis in CLL cells exposed to nucleoside analogues in vitro. *Eur J Haematol* **56**:213-20.
- Phillips KE and Munson AE (1997) 2'3'-Dideoxyinosine inhibits the humoral immune response in female B6C3F1 mice by targeting the B lymphocyte. *Toxicol Appl Pharmacol* **145**:260-7.
- Pinto-Duarte A, Coelho JE, Cunha RA, Ribeiro JA and Sebastiao AM (2005) Adenosine A2A receptors control the extracellular levels of adenosine through modulation of nucleoside transporters activity in the rat hippocampus. *J Neurochem* **93**:595-604.
- Playford RJ, Hanby AM, Gschmeissner S, Peiffer LP, Wright NA and McGarrity T (1996) The epidermal growth factor receptor (EGF-R) is present on the basolateral, but not the apical, surface of enterocytes in the human gastrointestinal tract. *Gut* **39**:262-6.
- Pressacco J, Mitrovski B, Erlichman C and Hedley DW (1995) Effects of thymidylate synthase inhibition on thymidine kinase activity and nucleoside transporter expression. *Cancer Res* **55**:1505-8.
- Pugmire MJ and Ealick SE (2002) Structural analyses reveal two distinct families of nucleoside phosphorylases. *Biochem J* **361**:1-25.

Q

- Quamme G, Pfeilschifter J and Murer H (1989) Parathyroid hormone inhibition of Na+/phosphate cotransport in OK cells: intracellular [Ca²⁺] as a second messenger. *Biochim Biophys Acta* **1013**:166-72.
- Quan R and Barness LA (1990) Do infants need nucleotide supplemented formula for optimal nutrition? *J Pediatr Gastroenterol Nutr* **11**:429-34.
- Quaroni A, Tian JQ, Goke M and Podolsky DK (1999) Glucocorticoids have pleiotropic effects on small intestinal crypt cells. *Am J Physiol* **277**:G1027-40.
- Quaroni A, Wands J, Trelstad RL and Isselbacher KJ (1979) Epithelioid cell cultures from rat small intestine. Characterization by morphologic and immunologic criteria. *J Cell Biol* **80**:248-65.

R

- Ranaldi G, Consalvo R, Sambuy Y and Scarino ML (2003) Permeability characteristics of parental and clonal human intestinal Caco-2 cell lines differentiated in serum-supplemented and serum-free media. *Toxicol In Vitro* **17**:761-7.
- Rauchwerger DR, Firby PS, Hedley DW and Moore MJ (2000) Equilibrative-sensitive nucleoside transporter and its role in gemcitabine sensitivity. *Cancer Res* **60**:6075-9.
- Ritzel MW, Ng AM, Yao SY, Graham K, Loewen SK, Smith KM, Hyde RJ, Karpinski E, Cass CE, Baldwin SA and Young JD (2001a) Recent molecular advances in studies of the concentrative Na⁺-dependent nucleoside transporter (CNT) family: identification and characterization of novel human and mouse proteins (hCNT3 and mCNT3) broadly selective for purine and pyrimidine nucleosides (system cib). *Mol Membr Biol* **18**:65-72.

- Ritzel MW, Ng AM, Yao SY, Graham K, Loewen SK, Smith KM, Ritzel RG, Mowles DA, Carpenter P, Chen XZ, Karpinski E, Hyde RJ, Baldwin SA, Cass CE and Young JD (2001b) Molecular identification and characterization of novel human and mouse concentrative Na⁺-nucleoside cotransporter proteins (hCNT3 and mCNT3) broadly selective for purine and pyrimidine nucleosides (system cib). *J Biol Chem* **276**:2914-27.
- Ritzel MW, Yao SY, Huang MY, Elliott JF, Cass CE and Young JD (1997) Molecular cloning and functional expression of cDNAs encoding a human Na⁺-nucleoside cotransporter (hCNT1). *Am J Physiol* **272**:C707-14.
- Ritzel MW, Yao SY, Ng AM, Mackey JR, Cass CE and Young JD (1998) Molecular cloning, functional expression and chromosomal localization of a cDNA encoding a human Na⁺/nucleoside cotransporter (hCNT2) selective for purine nucleosides and uridine. *Mol Membr Biol* **15**:203-11.
- Roepstorff C, Vistisen B, Donsmark M, Nielsen JN, Galbo H, Green KA, Hardie DG, Wojtaszewski JF, Richter EA and Kiens B (2004) Regulation of hormone-sensitive lipase activity and Ser563 and Ser565 phosphorylation in human skeletal muscle during exercise. *J Physiol* **560**:551-62.
- Rokutan K, Sakai A, Teramoto F, Kido Y, Shizuka F and Kishi K (1994) Epidermal growth factor-induced mitogen signals in cultured intestinal epithelial cells. *J Gastroenterol Suppl* **7**:59-62.
- Roman RM and Fitz JG (1999) Emerging roles of purinergic signaling in gastrointestinal epithelial secretion and hepatobiliary function. *Gastroenterology* **116**:964-79.
- Ruderman NB, Park H, Kaushik VK, Dean D, Constant S, Prentki M and Saha AK (2003) AMPK as a metabolic switch in rat muscle, liver and adipose tissue after exercise. *Acta Physiol Scand* **178**:435-42.
- Rudolph FB, Kulkarni AD, Schandle VB and Van Buren CT (1984) Involvement of dietary nucleotides in T lymphocyte function. *Adv Exp Med Biol* **165 Pt B**:175-8.
- Ruiz-Montasell B, Javier Casado F, Felipe A and Pastor-Anglada M (1992) Uridine transport in basolateral plasma membrane vesicles from rat liver. *J Membr Biol* **128**:227-33.
- Rutter GA, Da Silva Xavier G and Leclerc I (2003) Roles of 5'-AMP-activated protein kinase (AMPK) in mammalian glucose homeostasis. *Biochem J* **375**:1-16.
- Ryder JW, Chibalin AV and Zierath JR (2001) Intracellular mechanisms underlying increases in glucose uptake in response to insulin or exercise in skeletal muscle. *Acta Physiol Scand* **171**:249-57.

S

- Saadjian AY, Levy S, Franceschi F, Zouher I, Paganelli F and Guieu RP (2002) Role of endogenous adenosine as a modulator of syncope induced during tilt testing. *Circulation* **106**:569-74.
- Saitoh M, Nagai K, Nakagawa K, Yamamura T, Yamamoto S and Nishizaki T (2004) Adenosine induces apoptosis in the human gastric cancer cells via an intrinsic pathway relevant to activation of AMP-activated protein kinase. *Biochem Pharmacol* **67**:2005-11.
- Sakowicz M, Grden M and Pawelczyk T (2001) Expression level of adenosine kinase in rat tissues. Lack of phosphate effect on the enzyme activity. *Acta Biochim Pol* **48**:745-54.
- Salati LM, Gross CJ, Henderson LM and Savaiano DA (1984) Absorption and metabolism of adenine, adenosine-5'-monophosphate, adenosine and hypoxanthine by the isolated vascularly perfused rat small intestine. *J Nutr* **114**:753-60.
- Salloum RM, Stevens BR, Schultz GS and Souba WW (1993) Regulation of small intestinal glutamine transport by epidermal growth factor. *Surgery* **113**:552-9.
- Salt I, Celler JW, Hawley SA, Prescott A, Woods A, Carling D and Hardie DG (1998) AMP-activated protein kinase: greater AMP dependence, and preferential nuclear localization, of complexes containing the alpha2 isoform. *Biochem J* **334 (Pt 1)**:177-87.
- Salt IP, Connell JM and Gould GW (2000) 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR) inhibits insulin-stimulated glucose transport in 3T3-L1 adipocytes. *Diabetes* **49**:1649-56.
- Sanderson IR and He Y (1994) Nucleotide uptake and metabolism by intestinal epithelial cells. *J Nutr* **124**:131S-137S.

- Schaner ME, Wang J, Zevin S, Gerstin KM and Giacomini KM (1997) Transient expression of a purine-selective nucleoside transporter (SPNTint) in a human cell line (HeLa). *Pharm Res* **14**:1316-21.
- Scharrer E and Grenacher B (2001) Active intestinal absorption of nucleosides by Na⁺-dependent transport across the brush border membrane in cows. *J Dairy Sci* **84**:614-9.
- Schlomme E, Martin D and Meisel H (2000) Nucleosides and nucleotides: natural bioactive substances in milk and colostrum. *Br J Nutr* **84 Suppl 1**:S59-68.
- Schwarz SM, Ling SD, Hostetler B, Draper JP and Watkins JB (1984) Lipid composition and membrane fluidity in the small intestine of the developing rabbit. *Gastroenterology* **86**:1544-51.
- Scott JW, Norman DG, Hawley SA, Kontogiannis L and Hardie DG (2002) Protein kinase substrate recognition studied using the recombinant catalytic domain of AMP-activated protein kinase and a model substrate. *J Mol Biol* **317**:309-23.
- Seglen PO (1976) Preparation of isolated rat liver cells. *Methods Cell Biol* **13**:29-83.
- Serre V, Guy H, Liu X, Perverne B, Herve G and Evans D (1998) Allosteric regulation and substrate channeling in multifunctional pyrimidine biosynthetic complexes: analysis of isolated domains and yeast-mammalian chimeric proteins. *J Mol Biol* **281**:363-77.
- Sessa WC, Barber CM and Lynch KR (1993) Mutation of N-myristoylation site converts endothelial cell nitric oxide synthase from a membrane to a cytosolic protein. *Circ Res* **72**:921-4.
- Sgarrella F, Poddie FP, Meloni MA, Sciola L, Pippia P and Tozzi MG (1997) Channelling of deoxyribose moiety of exogenous DNA into carbohydrate metabolism: role of deoxyriboaldolase. *Comp Biochem Physiol B Biochem Mol Biol* **117**:253-7.
- Shen F, Look KY, Yeh YA and Weber G (1998) Increased uridine kinase (ATP: uridine 5'-phosphotransferase; EC 2.7.1.48) activity in human and rat tumors. *Cancer Biochem Biophys* **16**:1-15.
- Shepherd EJ, Hellwell PA, Mace OJ, Morgan EL, Patel N and Kellett GL (2004) Stress and glucocorticoid inhibit apical GLUT2-trafficking and intestinal glucose absorption in rat small intestine. *J Physiol* **560**:281-90.
- Shewach DS, Ellero J, Mancini WR and Ensminger WD (1992) Decrease in TTP pools mediated by 5-bromo-2'-deoxyuridine exposure in a human glioblastoma cell line. *Biochem Pharmacol* **43**:1579-85.
- Shi MM, Wu JS, Lee CM and Young JD (1984) Nucleoside transport. Photoaffinity labelling of high-affinity nitrobenzylthioinosine binding sites in rat and guinea pig lung. *Biochem Biophys Res Commun* **118**:594-600.
- Shin HC, Landowski CP, Sun D, Vig BS, Kim I, Mittal S, Lane M, Rosania G, Drach JC and Amidon GL (2003) Functional expression and characterization of a sodium-dependent nucleoside transporter hCNT2 cloned from human duodenum. *Biochem Biophys Res Commun* **307**:696-703.
- Shiraga T, Miyamoto K, Tanaka H, Yamamoto H, Taketani Y, Morita K, Tamai I, Tsuji A and Takeda E (1999) Cellular and molecular mechanisms of dietary regulation on rat intestinal H⁺/Peptide transporter PepT1. *Gastroenterology* **116**:354-62.
- Sigoillot FD, Berkowski JA, Sigoillot SM, Kotsis DH and Guy HI (2003) Cell cycle-dependent regulation of pyrimidine biosynthesis. *J Biol Chem* **278**:3403-9.
- Sigoillot FD, Evans DR and Guy HI (2002) Growth-dependent regulation of mammalian pyrimidine biosynthesis by the protein kinase A and MAPK signaling cascades. *J Biol Chem* **277**:15745-51.
- Snyder FF and Lukey T (1982) Kinetic considerations for the regulation of adenosine and deoxyadenosine metabolism in mouse and human tissues based on a thymocyte model. *Biochim Biophys Acta* **696**:299-307.
- Sobrero AF, Moir RD, Bertino JR and Handschumacher RE (1985) Defective facilitated diffusion of nucleosides, a primary mechanism of resistance to 5-fluoro-2'-deoxyuridine in the HCT-8 human carcinoma line. *Cancer Res* **45**:3155-60.
- Soler C, Felipe A, Casado FJ, Celada A and Pastor-Anglada M (2000) Nitric oxide regulates nucleoside transport in activated B lymphocytes. *J Leukoc Biol* **67**:345-9.
- Soler C, Felipe A, Mata JF, Casado FJ, Celada A and Pastor-Anglada M (1998) Regulation of nucleoside transport by lipopolysaccharide, phorbol esters, and tumor necrosis factor-alpha in human B-lymphocytes. *J Biol Chem* **273**:26939-45.

- Soler C, Garcia-Manteiga J, Valdes R, Xaus J, Comalada M, Casado FJ, Pastor-Anglada M, Celada A and Felipe A (2001a) Macrophages require different nucleoside transport systems for proliferation and activation. *Faseb J* **15**:1979-88.
- Soler C, Valdes R, Garcia-Manteiga J, Xaus J, Comalada M, Casado FJ, Modolell M, Nicholson B, MacLeod C, Felipe A, Celada A and Pastor-Anglada M (2001b) Lipopolysaccharide-induced apoptosis of macrophages determines the up-regulation of concentrative nucleoside transporters Cnt1 and Cnt2 through tumor necrosis factor-alpha-dependent and -independent mechanisms. *J Biol Chem* **276**:30043-9.
- Somasekaram A, Jarmuz A, How A, Scott J and Navaratnam N (1999) Intracellular localization of human cytidine deaminase. Identification of a functional nuclear localization signal. *J Biol Chem* **274**:28405-12.
- Spiegelman D, Carroll RJ and Kipnis V (2001) Efficient regression calibration for logistic regression in main study/internal validation study designs with an imperfect reference instrument. *Stat Med* **20**:139-160.
- Stein SC, Woods A, Jones NA, Davison MD and Carling D (2000) The regulation of AMP-activated protein kinase by phosphorylation. *Biochem J* **345 Pt 3**:437-43.
- Stiles GL (1992) Adenosine receptors. *J Biol Chem* **267**:6451-4.
- Stolk M, Cooper E, Vilk G, Litchfield DW and Hammond JR (2005) Subtype-specific regulation of equilibrative nucleoside transporters by protein kinase CK2. *Biochem J* **386**:281-9.
- Suemori S, Ciacci C and Podolsky DK (1991) Regulation of transforming growth factor expression in rat intestinal epithelial cell lines. *J Clin Invest* **87**:2216-21.
- Sugiura M, Kono K, Liu H, Shimizugawa T, Minekura H, Spiegel S and Kohama T (2002) Ceramide kinase, a novel lipid kinase. Molecular cloning and functional characterization. *J Biol Chem* **277**:23294-300.
- Sun AQ, Ananthanarayanan M, Soroka CJ, Thevananther S, Shneider BL and Suchy FJ (1998) Sorting of rat liver and ileal sodium-dependent bile acid transporters in polarized epithelial cells. *Am J Physiol* **275**:G1045-55.
- Sun AQ, Balasubramanyan N, Liu CJ, Shahid M and Suchy FJ (2004) Association of the 16-kDa subunit c of vacuolar proton pump with the ileal Na+-dependent bile acid transporter: protein-protein interaction and intracellular trafficking. *J Biol Chem* **279**:16295-300.
- Sun AQ, Salkar R, Sachchidanand, Xu S, Zeng L, Zhou MM and Suchy FJ (2003) A 14-amino acid sequence with a beta-turn structure is required for apical membrane sorting of the rat ileal bile acid transporter. *J Biol Chem* **278**:4000-9.
- Sundaram M, Yao SY, Ingram JC, Berry ZA, Abidi F, Cass CE, Baldwin SA and Young JD (2001) Topology of a human equilibrative, nitrobenzylthioinosine (NBMPR)-sensitive nucleoside transporter (hENT1) implicated in the cellular uptake of adenosine and anti-cancer drugs. *J Biol Chem* **276**:45270-5.
- Sundaram M, Yao SY, Ng AM, Griffiths M, Cass CE, Baldwin SA and Young JD (1998) Chimeric constructs between human and rat equilibrative nucleoside transporters (hENT1 and rENT1) reveal hENT1 structural domains interacting with coronary vasoactive drugs. *J Biol Chem* **273**:21519-25.
- Szkotak AJ, Man SF and Duszyk M (2003) The role of the basolateral outwardly rectifying chloride channel in human airway epithelial anion secretion. *Am J Respir Cell Mol Biol* **29**:710-20.
- Szkotak AJ, Ng AM, Sawicka J, Baldwin SA, Man SF, Cass CE, Young JD and Duszyk M (2001) Regulation of K(+) current in human airway epithelial cells by exogenous and autocrine adenosine. *Am J Physiol Cell Physiol* **281**:C1991-2002.

T

- Tattersall MH, Slowiaczek P and De Fazio A (1983) Regional variation in human extracellular purine levels. *J Lab Clin Med* **102**:411-20.
- Thiesen A, Drozdowski L, lordache C, Neo CC, Woudstra TD, Xenodemetropoulos T, Keelan M, Clandinin MT, Thomson AB and Wild G (2003) Adaptation following intestinal resection: mechanisms and signals. *Best Pract Res Clin Gastroenterol* **17**:981-95.
- Thwaites DT, Brown CD, Hirst BH and Simmons NL (1993) H(+) -coupled dipeptide (glycylsarcosine) transport across apical and basal borders of human intestinal Caco-2 cell monolayers display distinctive characteristics. *Biochim Biophys Acta* **1151**:237-45.

- Trimble ME and Coulson R (1984) Adenosine transport in perfused rat kidney and renal cortical membrane vesicles. *Am J Physiol* **246**:F794-803.
- Tse CM, Belt JA, Jarvis SM, Paterson AR, Wu JS and Young JD (1985) Reconstitution studies of the human erythrocyte nucleoside transporter. *J Biol Chem* **260**:3506-11.

V

- Vaandrager AB, Ehlert EM, Jarchau T, Lohmann SM and de Jonge HR (1996) N-terminal myristylation is required for membrane localization of cGMP-dependent protein kinase type II. *J Biol Chem* **271**:7025-9.
- Valdes R, Casado FJ and Pastor-Anglada M (2002) Cell-cycle-dependent regulation of CNT1, a concentrative nucleoside transporter involved in the uptake of cell-cycle-dependent nucleoside-derived anticancer drugs. *Biochem Biophys Res Commun* **296**:575-9.
- Valdes R, Ortega MA, Casado FJ, Felipe A, Gil A, Sanchez-Pozo A and Pastor-Anglada M (2000) Nutritional regulation of nucleoside transporter expression in rat small intestine. *Gastroenterology* **119**:1623-30.
- Van Rompay AR, Johansson M and Karlsson A (1999) Phosphorylation of deoxycytidine analog monophosphates by UMP-CMP kinase: molecular characterization of the human enzyme. *Mol Pharmacol* **56**:562-9.
- Van Rompay AR, Johansson M and Karlsson A (2000) Phosphorylation of nucleosides and nucleoside analogs by mammalian nucleoside monophosphate kinases. *Pharmacol Ther* **87**:189-98.
- Van Rompay AR, Norda A, Linden K, Johansson M and Karlsson A (2001) Phosphorylation of uridine and cytidine nucleoside analogs by two human uridine-cytidine kinases. *Mol Pharmacol* **59**:1181-6.
- Vavvas D, Apazidis A, Saha AK, Gamble J, Patel A, Kemp BE, Witters LA and Ruderman NB (1997) Contraction-induced changes in acetyl-CoA carboxylase and 5'-AMP-activated kinase in skeletal muscle. *J Biol Chem* **272**:13255-61.
- Vickers MF, Mani RS, Sundaram M, Hogue DL, Young JD, Baldwin SA and Cass CE (1999) Functional production and reconstitution of the human equilibrative nucleoside transporter (hENT1) in *Saccharomyces cerevisiae*. Interaction of inhibitors of nucleoside transport with recombinant hENT1 and a glycosylation-defective derivative (hENT1/N48Q). *Biochem J* **339 (Pt 1)**:21-32.
- Vijayalakshmi D and Belt JA (1988) Sodium-dependent nucleoside transport in mouse intestinal epithelial cells. Two transport systems with differing substrate specificities. *J Biol Chem* **263**:19419-23.
- Vijayalakshmi D, Dagnino L, Belt JA, Gati WP, Cass CE and Paterson AR (1992) L1210/B23.1 cells express equilibrative, inhibitor-sensitive nucleoside transport activity and lack two parental nucleoside transport activities. *J Biol Chem* **267**:16951-6.

W

- Wang J and Giacomini KM (1999a) Characterization of a bioengineered chimeric Na⁺-nucleoside transporter. *Mol Pharmacol* **55**:234-40.
- Wang J and Giacomini KM (1999b) Serine 318 is essential for the pyrimidine selectivity of the N2 Na⁺-nucleoside transporter. *J Biol Chem* **274**:2298-302.
- Wang J, Su SF, Dresser MJ, Schaner ME, Washington CB and Giacomini KM (1997) Na(+)-dependent purine nucleoside transporter from human kidney: cloning and functional characterization. *Am J Physiol* **273**:F1058-65.
- Wang Y and Oram JF (2005) Unsaturated fatty acids phosphorylate and destabilize ABCA1 through a phospholipase D2 pathway. *J Biol Chem*.
- Ward JL, Leung GP, Toan SV and Tse CM (2003) Functional analysis of site-directed glycosylation mutants of the human equilibrative nucleoside transporter-2. *Arch Biochem Biophys* **411**:19-26.
- Ward JL, Sherli A, Mo ZP and Tse CM (2000) Kinetic and pharmacological properties of cloned human equilibrative nucleoside transporters, ENT1 and ENT2, stably expressed in nucleoside transporter-deficient PK15 cells. Ent2 exhibits a low affinity for guanosine and cytidine but a high affinity for inosine. *J Biol Chem* **275**:8375-81.

- Ward JL and Tse CM (1999) Nucleoside transport in human colonic epithelial cell lines: evidence for two Na⁺-independent transport systems in T84 and Caco-2 cells. *Biochim Biophys Acta* **1419**:15-22.
- Weber G (1983) Enzymes of purine metabolism in cancer. *Clin Biochem* **16**:57-63.
- Weber G, Nakamura H, Natsumeda Y, Szekeres T and Nagai M (1992) Regulation of GTP biosynthesis. *Adv Enzyme Regul* **32**:57-69.
- Wheeler EE and Challacombe DN (1997) The trophic action of growth hormone, insulin-like growth factor-I, and insulin on human duodenal mucosa cultured in vitro. *Gut* **40**:57-60.
- Wiley JS, Cebon JS, Jamieson GP, Szer J, Gibson J, Woodruff RK, McKendrick JJ, Sheridan WP, Biggs JC, Snook MB and et al. (1994) Assessment of proliferative responses to granulocyte-macrophage colony-stimulating factor (GM-CSF) in acute myeloid leukaemia using a fluorescent ligand for the nucleoside transporter. *Leukemia* **8**:181-5.
- Wiley JS, Jones SP, Sawyer WH and Paterson AR (1982) Cytosine arabinoside influx and nucleoside transport sites in acute leukemia. *J Clin Invest* **69**:479-89.
- Williams TC, Doherty AJ, Griffith DA and Jarvis SM (1989) Characterization of sodium-dependent and sodium-independent nucleoside transport systems in rabbit brush-border and basolateral plasma-membrane vesicles from the renal outer cortex. *Biochem J* **264**:223-31.
- Williams TC and Jarvis SM (1990) Na(+) -dependent purine and pyrimidine nucleoside transporters in bovine outer renal cortex brush-border-membrane vesicles. *Biochem Soc Trans* **18**:684-5.
- Williams TC and Jarvis SM (1991) Multiple sodium-dependent nucleoside transport systems in bovine renal brush-border membrane vesicles. *Biochem J* **274 (Pt 1)**:27-33.
- Wilmore DW and Robinson MK (2000) Short bowel syndrome. *World J Surg* **24**:1486-92.
- Winder WW and Hardie DG (1999) AMP-activated protein kinase, a metabolic master switch: possible roles in type 2 diabetes. *Am J Physiol* **277**:E1-10.
- Wojtaszewski JF, Jorgensen SB, Hellsten Y, Hardie DG and Richter EA (2002) Glycogen-dependent effects of 5-aminoimidazole-4-carboxamide (AICA)-riboside on AMP-activated protein kinase and glycogen synthase activities in rat skeletal muscle. *Diabetes* **51**:284-92.
- Woods A, Azzout-Marniche D, Foretz M, Stein SC, Lemarchand P, Ferre P, Foufelle F and Carling D (2000) Characterization of the role of AMP-activated protein kinase in the regulation of glucose-activated gene expression using constitutively active and dominant negative forms of the kinase. *Mol Cell Biol* **20**:6704-11.
- Woods A, Johnstone SR, Dickerson K, Leiper FC, Fryer LG, Neumann D, Schlattner U, Wallimann T, Carlson M and Carling D (2003) LKB1 is the upstream kinase in the AMP-activated protein kinase cascade. *Curr Biol* **13**:2004-8.
- Wright AM, Paterson AR, Sowa B, Akabutu JJ, Grundy PE and Gati WP (2002) Cytotoxicity of 2-chlorodeoxyadenosine and arabinosylcytosine in leukaemic lymphoblasts from paediatric patients: significance of cellular nucleoside transporter content. *Br J Haematol* **116**:528-37.
- Wright EM (2001) Renal Na(+) -glucose cotransporters. *Am J Physiol Renal Physiol* **280**: F10-8.
- Wright EM, Martin MG and Turk E (2003) Intestinal absorption in health and disease--sugars. *Best Pract Res Clin Gastroenterol* **17**:943-56.
- Wu JS and Young JD (1984) Photoaffinity labelling of nucleoside-transport proteins in plasma membranes isolated from rat and guinea-pig liver. *Biochem J* **220**:499-506.
- Wu X, Gutierrez MM and Giacomini KM (1994) Further characterization of the sodium-dependent nucleoside transporter (N3) in choroid plexus from rabbit. *Biochim Biophys Acta* **1191**:190-6.
- Wu X, Motoshima H, Mahadev K, Stalker TJ, Scalia R and Goldstein BJ (2003) Involvement of AMP-activated protein kinase in glucose uptake stimulated by the globular domain of adiponectin in primary rat adipocytes. *Diabetes* **52**:1355-63.

X

- Xiao G, Wang J, Tangen T and Giacomini KM (2001) A novel proton-dependent nucleoside transporter, CeCNT3, from *Caenorhabditis elegans*. *Mol Pharmacol* **59**:339-48.
- Xu PA and Kellem RE (2000) Function of murine adenosine deaminase in the gastrointestinal tract. *Biochem Biophys Res Commun* **269**:749-57.

Y

- Yamaguchi N, Kodama M and Ueda K (1985) Diadenosine tetraphosphate as a signal molecule linked with the functional state of rat liver. *Gastroenterology* **89**:723-31.
- Yamaoka T and Itakura M (1996) [Metabolism of purine nucleotides and the production of uric acid]. *Nippon Rinsho* **54**:3188-94.
- Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, Yamashita S, Noda M, Kita S, Ueki K, Eto K, Akanuma Y, Froguel P, Foufelle F, Ferre P, Carling D, Kimura S, Nagai R, Kahn BB and Kadokawa T (2002) Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* **8**:1288-95.
- Yao SY, Cass CE and Young JD (1996) Transport of the antiviral nucleoside analogs 3'-azido-3'-deoxythymidine and 2',3'-dideoxycytidine by a recombinant nucleoside transporter (rCNT) expressed in Xenopus laevis oocytes. *Mol Pharmacol* **50**:388-93.
- Yao SY, Ng AM, Loewen SK, Cass CE, Baldwin SA and Young JD (2002a) An ancient prevertebrate Na⁺-nucleoside cotransporter (hfCNT) from the Pacific hagfish (*Eptatretus stouti*). *Am J Physiol Cell Physiol* **283**:C155-68.
- Yao SY, Ng AM, Muzyka WR, Griffiths M, Cass CE, Baldwin SA and Young JD (1997) Molecular cloning and functional characterization of nitrobenzylthioinosine (NBMPR)-sensitive (es) and NBMPR-insensitive (ei) equilibrative nucleoside transporter proteins (rENT1 and rENT2) from rat tissues. *J Biol Chem* **272**:28423-30.
- Yao SY, Ng AM, Sundaram M, Cass CE, Baldwin SA and Young JD (2001a) Transport of antiviral 3'-deoxy-nucleoside drugs by recombinant human and rat equilibrative, nitrobenzylthioinosine (NBMPR)-insensitive (ENT2) nucleoside transporter proteins produced in Xenopus oocytes. *Mol Membr Biol* **18**:161-7.
- Yao SY, Ng AM, Vickers MF, Sundaram M, Cass CE, Baldwin SA and Young JD (2002b) Functional and molecular characterization of nucleobase transport by recombinant human and rat equilibrative nucleoside transporters 1 and 2. Chimeric constructs reveal a role for the ENT2 helix 5-6 region in nucleobase translocation. *J Biol Chem* **277**:24938-48.
- Yao SY, Sundaram M, Chomey EG, Cass CE, Baldwin SA and Young JD (2001b) Identification of Cys140 in helix 4 as an exofacial cysteine residue within the substrate-translocation channel of rat equilibrative nitrobenzylthioinosine (NBMPR)-insensitive nucleoside transporter rENT2. *Biochem J* **353**:387-93.
- Yasuda K, Kosugi A, Hayashi F, Saitoh S, Nagafuku M, Mori Y, Ogata M and Hamaoka T (2000) Serine 6 of Lck tyrosine kinase: a critical site for Lck myristylation, membrane localization, and function in T lymphocytes. *J Immunol* **165**:3226-31.

Z

- Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, Wu M, Ventre J, Doepper T, Fujii N, Musi N, Hirshman MF, Goodyear LJ and Moller DE (2001) Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest* **108**:1167-74.
- Zimmermann H (2000) Extracellular metabolism of ATP and other nucleotides. *Naunyn Schmiedebergs Arch Pharmacol* **362**:299-309.