

Programa de Doctorat de Biomedicina
Departament de Bioquímica i Biologia Molecular
Facultat de Biologia, Universitat de Barcelona
Bienni 2005-2007

NOVES FUNCIONS DE LA PROTEÏNA FLOTILLIN-1 EN LA REGULACIÓ DEL PROCÉS DE MITOSI I DE LA VIA DE SENYALITZACIÓ DEL RECEPTOR NOTCH1

Memòria presentada per
Valentí Gómez Martínez

per optar al grau de
Doctor
per la Universitat de Barcelona

Treball realitzat sota la direcció de la Dra. Rosanna Paciucci, a la Unitat de Recerca Biomèdica de l’Institut de Recerca de l’Hospital Universitari Vall d’Hebron.
Amb el suport del Departament d’Universitats, Recerca i Societat de la Informació de la Generalitat de Catalunya.

Tesi doctoral adscrita al departament de Bioquímica i Biologia Molecular de la Facultat de Biologia, Universitat de Barcelona, sota la tutoria de la Dra. Adela Mazo.

Dra. Rosanna Paciucci
directora de tesi

Dra. Adela Mazo
tutora de tesi

Valentí Gómez
autor

Barcelona, Abril de 2009

BIBLIOGRAFIA



BIBLIOGRAFIA

A

Aagaard L, Rossi JJ (2007) RNAi therapeutics: principles, prospects and challenges. *Adv Drug Deliv Rev* 59(2-3): 75-86

Abdullah AS, Foong C, Murata-Hori M (2005) Specific distribution of overexpressed Aurora B kinase during interphase of normal epithelial cells. *Cancer Cell Int* 5: 31

Adams RR, Maiato H, Earnshaw WC, Carmena M (2001) Essential roles of Drosophila inner centromere protein (INCENP) and Aurora B in histone H3 phosphorylation, metaphase chromosome alignment, kinetochore disjunction, and chromosome segregation. *J Cell Biol* 153(4): 865-880

Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P (2002) *Molecular Biology of the Cell*, 4th Edition edn.

Albertson R, Cao J, Hsieh TS, Sullivan W (2008) Vesicles and actin are targeted to the cleavage furrow via furrow microtubules and the central spindle. *J Cell Biol* 181(5): 777-790

Allman D, Punt JA, Izon DJ, Aster JC, Pear WS (2002) An invitation to T and more: notch signaling in lymphopoiesis. *Cell* 109 Suppl: S1-11

Andreassen PR, Lacroix FB, Lohez OD, Margolis RL (2001) Neither p21WAF1 nor 14-3-3sigma prevents G2 progression to mitotic catastrophe in human colon carcinoma cells after DNA damage, but p21WAF1 induces stable G1 arrest in resulting tetraploid cells. *Cancer Res* 61(20): 7660-7668

Artavanis-Tsakonas S, Matsuno K, Fortini ME (1995) Notch signaling. *Science* 268(5208): 225-232

Artavanis-Tsakonas S, Rand MD, Lake RJ (1999) Notch signaling: cell fate control and signal integration in development. *Science* 284(5415): 770-776

B

Baumann CA, Ribon V, Kanzaki M, Thurmond DC, Mora S, Shigematsu S, Bickel PE, Pessin JE, Saltiel AR (2000) CAP defines a second signalling pathway required for insulin-stimulated glucose transport. *Nature* 407(6801): 202-207

Benson LJ, Gu Y, Yakovleva T, Tong K, Barrows C, Strack CL, Cook RG, Mizzen CA, Annunziato AT (2006) Modifications of H3 and H4 during chromatin replication, nucleosome assembly, and histone exchange. *J Biol Chem* 281(14): 9287-9296

Berdnik D, Knoblich JA (2002) Drosophila Aurora-A is required for centrosome maturation and actin-dependent asymmetric protein localization during mitosis. *Curr Biol* 12(8): 640-647

Berdnik D, Torok T, Gonzalez-Gaitan M, Knoblich JA (2002) The endocytic protein alpha-Adaptin is required for numb-mediated asymmetric cell division in Drosophila. *Dev Cell* 3(2): 221-231

Bickel PE, Scherer PE, Schnitzer JE, Oh P, Lisanti MP, Lodish HF (1997) Flotillin and epidermal surface antigen define a new family of caveolae-associated integral membrane proteins. *J Biol Chem* 272(21): 13793-13802

Blasina A, de Weyer IV, Laus MC, Luyten WH, Parker AE, McGowan CH (1999) A human homologue of the checkpoint kinase Cds1 directly inhibits Cdc25 phosphatase. *Curr Biol* 9(1): 1-10

Blaumueller CM, Qi H, Zagouras P, Artavanis-Tsakonas S (1997) Intracellular cleavage of Notch leads to a heterodimeric receptor on the plasma membrane. *Cell* 90(2): 281-291

Brou C, Logeat F, Gupta N, Bessia C, LeBail O, Doedens JR, Cumano A, Roux P, Black RA, Israel A (2000) A novel proteolytic cleavage involved in Notch signaling: the role of the disintegrin-metalloprotease TACE. *Mol Cell* 5(2): 207-216

Browman DT, Hoegg MB, Robbins SM (2007) The SPFH domain-containing proteins: more than lipid raft markers. *Trends Cell Biol* 17(8): 394-402

Brown DA, Rose JK (1992) Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface. *Cell* 68(3): 533-544

Burke R, Nellen D, Bellotto M, Hafen E, Senti KA, Dickson BJ, Basler K (1999) Dispatched, a novel sterol-sensing domain protein dedicated to the release of cholesterol-modified hedgehog from signaling cells. *Cell* 99(7): 803-815

Bush G, diSibio G, Miyamoto A, Denault JB, Leduc R, Weinmaster G (2001) Ligand-induced signaling in the absence of furin processing of Notch1. *Dev Biol* 229(2): 494-502

C

Campos LS, Decker L, Taylor V, Skarnes W (2006) Notch, epidermal growth factor receptor, and beta1-integrin pathways are coordinated in neural stem cells. *J Biol Chem* 281(8): 5300-5309

Carmena M, Earnshaw WC (2003) The cellular geography of Aurora kinases. *Nat Rev Mol Cell Biol* 4(11): 842-854

Carvalho A, Carmena M, Sambade C, Earnshaw WC, Wheatley SP (2003) Survivin is required for stable checkpoint activation in taxol-treated HeLa cells. *J Cell Sci* 116(Pt 14): 2987-2998

Castro A, Arlot-Bonnemains Y, Vigneron S, Labbe JC, Prigent C, Lorca T (2002) APC/Fizzy-Related targets Aurora-A kinase for proteolysis. *EMBO Rep* 3(5): 457-462

Chan HM, La Thangue NB (2001) p300/CBP proteins: HATs for transcriptional bridges and scaffolds. *J Cell Sci* 114(Pt 13): 2363-2373

Chen JD, Evans RM (1995) A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* 377(6548): 454-457

Chen TY, Liu PH, Ruan CT, Chiu L, Kung FL (2006) The intracellular domain of amyloid precursor protein interacts with Flotillin-1, a lipid raft protein. *Biochem Biophys Res Commun* 342(1): 266-272

Cheng H, Vetrikel KS, Drisdel RC, Meckler X, Gong P, Leem JY, Li T, Carter M, Chen Y, Nguyen P, Iwatsubo T, Tomita T, Wong PC, Green WN, Kounnas MZ, Thinakaran G (2009) S-Palmitoylation of {gamma}-Secretase Subunits Nicastrin and APH-1. *J Biol Chem* 284(3): 1373-1384

Chomczynski P, Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162(1): 156-159

Chow JP, Siu WY, Fung TK, Chan WM, Lau A, Arooz T, Ng CP, Yamashita K, Poon RY (2003) DNA damage during the spindle-assembly checkpoint degrades CDC25A, inhibits cyclin-CDC2 complexes, and reverses cells to interphase. *Mol Biol Cell* 14(10): 3989-4002

Conner SD, Schmid SL (2003) Regulated portals of entry into the cell. *Nature* 422(6927): 37-44

D

Dai J, Sultan S, Taylor SS, Higgins JM (2005) The kinase haspin is required for mitotic histone H3 Thr 3 phosphorylation and normal metaphase chromosome alignment. *Genes Dev* 19(4): 472-488

De Strooper B, Annaert W, Cupers P, Saftig P, Craessaerts K, Mumm JS, Schroeter EH, Schrijvers V, Wolfe MS, Ray WJ, Goate A, Kopan R (1999) A presenilin-1-dependent gamma-secretase-like protease mediates release of Notch intracellular domain. *Nature* 398(6727): 518-522

Descombes P, Nigg EA (1998) The polo-like kinase Plx1 is required for M phase exit and destruction of mitotic regulators in Xenopus egg extracts. *EMBO J* 17(5): 1328-1335

Ditchfield C, Johnson VL, Tighe A, Ellston R, Haworth C, Johnson T, Mortlock A, Keen N, Taylor SS (2003) Aurora B couples chromosome alignment with anaphase by targeting BubR1, Mad2, and Cenp-E to kinetochores. *J Cell Biol* 161(2): 267-280

Doherty SD, Prieto VG, George S, Hazarika P, Duvic M (2006) High Flotillin-2 expression is associated with lymph node metastasis and Breslow depth in melanoma. *Melanoma Res* 16(5): 461-463

Drab M, Verkade P, Elger M, Kasper M, Lohn M, Lauterbach B, Menne J, Lindschau C, Mende F, Luft FC, Schedl A, Haller H, Kurzchalia TV (2001) Loss of caveolae, vascular dysfunction, and pulmonary defects in Caveolin-1 gene-disrupted mice. *Science* 293(5539): 2449-2452

E

Edgar AJ, Polak JM (2001) Flotillin-1: gene structure: cDNA cloning from human lung and the identification of alternative polyadenylation signals. *Int J Biochem Cell Biol* 33(1): 53-64

Ellisen LW, Bird J, West DC, Soreng AL, Reynolds TC, Smith SD, Sklar J (1991) TAN-1, the human homolog of the Drosophila notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms. *Cell* 66(4): 649-661

F

Fanning AS, Anderson JM (1999) PDZ domains: fundamental building blocks in the organization of protein complexes at the plasma membrane. *J Clin Invest* 103(6): 767-772

Fernandez I, Ying Y, Albanesi J, Anderson RG (2002) Mechanism of Caveolin filament assembly. *Proc Natl Acad Sci U S A* 99(17): 11193-11198

Field KA, Holowka D, Baird B (1995) Fc epsilon RI-mediated recruitment of p53/56lyn to detergent-resistant membrane domains accompanies cellular signaling. *Proc Natl Acad Sci U S A* 92(20): 9201-9205

Fitzgerald K, Harrington A, Leder P (2000) Ras pathway signals are required for notch-mediated oncogenesis. *Oncogene* 19(37): 4191-4198

Fluza UM, Arias AM (2007) Cell and molecular biology of Notch. *J Endocrinol* 194(3): 459-474

Fleming RJ (1998) Structural conservation of Notch receptors and ligands. *Semin Cell Dev Biol* 9(6): 599-607

Frick M, Bright NA, Riento K, Bray A, Merrified C, Nichols BJ (2007) Coassembly of Flotillins induces formation of membrane microdomains, membrane curvature, and vesicle budding. *Curr Biol* 17(13): 1151-1156

Fuller BG, Lampson MA, Foley EA, Rosasco-Nitcher S, Le KV, Tobelmann P, Brautigan DL, Stukenberg PT, Kapoor TM (2008) Midzone activation of Aurora B in anaphase produces an intracellular phosphorylation gradient. *Nature* 453(7198): 1132-1136

Fuwa TJ, Hori K, Sasamura T, Higgs J, Baron M, Matsuno K (2006) The first deltex null mutant indicates tissue-specific deltex-dependent Notch signaling in Drosophila. *Mol Genet Genomics* 275(3): 251-263

G

Gamble SC, Chotai D, Odontiadis M, Dart DA, Brooke GN, Powell SM, Reebye V, Varela-Carver A, Kawano Y, Waxman J, Bevan CL (2007) Prohibitin, a protein downregulated by androgens, represses androgen receptor activity. *Oncogene* 26(12): 1757-1768

Garin J, Diez R, Kieffer S, Dermine JF, Duclos S, Gagnon E, Sadoul R, Rondeau C, Desjardins M (2001) The phagosome proteome: insight into phagosome functions. *J Cell Biol* 152(1): 165-180

Gassmann R, Carvalho A, Henzing AJ, Ruchaud S, Hudson DF, Honda R, Nigg EA, Gerloff DL, Earnshaw WC (2004) Borealin: a novel chromosomal passenger required for stability of the bipolar mitotic spindle. *J Cell Biol* 166(2): 179-191

Giot L, Bader JS, Brouwer C, Chaudhuri A, Kuang B, Li Y, Hao YL, Ooi CE, Godwin B, Vitols E, Vijayadamodar G, Pochart P, Machineni H, Welsh M, Kong Y, Zerhusen B, Malcolm R, Varrone Z, Collis A, Minto M, Burgess S, McDaniel L, Stimpson E, Spriggs F, Williams J, Neurath K, Ioime N, Agee M, Voss E, Furtak K, Renzulli R, Aanensen N, Carrolla S, Bickelhaupt E, Lazovatsky Y, DaSilva A, Zhong J, Stanyon CA, Finley RL, Jr., White KP, Braverman M, Jarvie T, Gold S, Leach M, Knight J, Shimkets RA, McKenna MP, Chant J, Rothberg JM (2003) A protein interaction map of *Drosophila melanogaster*. *Science* 302(5651): 1727-1736

Gkantiragas I, Brugger B, Stuven E, Kaloyanova D, Li XY, Lohr K, Lottspeich F, Wieland FT, Helms JB (2001) Sphingomyelin-enriched microdomains at the Golgi complex. *Mol Biol Cell* 12(6): 1819-1833

Glebov OO, Bright NA, Nichols BJ (2006) Flotillin-1 defines a clathrin-independent endocytic pathway in mammalian cells. *Nat Cell Biol* 8(1): 46-54

Glotzer M, Murray AW, Kirschner MW (1991) Cyclin is degraded by the ubiquitin pathway. *Nature* 349(6305): 132-138

Grossberger R, Gieffers C, Zachariae W, Podtelejnikov AV, Schleiffer A, Nasmyth K, Mann M, Peters JM (1999) Characterization of the DOC1/APC10 subunit of the yeast and the human anaphase-promoting complex. *J Biol Chem* 274(20): 14500-14507

Gupta-Rossi N, Six E, LeBail O, Logeat F, Chastagner P, Olry A, Israel A, Brou C (2004) Monoubiquitination and endocytosis direct gamma-secretase cleavage of activated Notch receptor. *J Cell Biol* 166(1): 73-83

H

Harper JW, Burton JL, Solomon MJ (2002) The anaphase-promoting complex: it's not just for mitosis any more. *Genes Dev* 16(17): 2179-2206

Harris BZ, Lim WA (2001) Mechanism and role of PDZ domains in signaling complex assembly. *J Cell Sci* 114(Pt 18): 3219-3231

Hauf S, Cole RW, LaTerra S, Zimmer C, Schnapp G, Walter R, Heckel A, van Meel J, Rieder CL, Peters JM (2003) The small molecule Hesperadin reveals a role for Aurora B in correcting kinetochore-microtubule attachment and in maintaining the spindle assembly checkpoint. *J Cell Biol* 161(2): 281-294

Hauf S, Waizenegger IC, Peters JM (2001) Cohesin cleavage by separase required for anaphase and cytokinesis in human cells. *Science* 293(5533): 1320-1323

Hayward P, Brennan K, Sanders P, Balayo T, DasGupta R, Perrimon N, Martinez Arias A (2005) Notch modulates Wnt signalling by associating with Armadillo/beta-catenin and regulating its transcriptional activity. *Development* 132(8): 1819-1830

Hazarika P, McCarty MF, Prieto VG, George S, Babu D, Koul D, Bar-Eli M, Duvic M (2004) Up-regulation of Flotillin-2 is associated with melanoma progression and modulates expression of the thrombin receptor protease activated receptor 1. *Cancer Res* 64(20): 7361-7369

Heitzler P, Simpson P (1993) Altered epidermal growth factor-like sequences provide evidence for a role of Notch as a receptor in cell fate decisions. *Development* 117(3): 1113-1123

Hinshaw JE (2000) Dynamin and its role in membrane fission. *Annu Rev Cell Dev Biol* 16: 483-519

Hirano T (2000) Chromosome cohesion, condensation, and separation. *Annu Rev Biochem* 69: 115-144

Hodkinson PS, Elliott PA, Lad Y, McHugh BJ, MacKinnon AC, Haslett C, Sethi T (2007) Mammalian NOTCH-1 activates beta1 integrins via the small GTPase R-Ras. *J Biol Chem* 282(39): 28991-29001

Hoehne M, de Couet HG, Stuermer CA, Fischbach KF (2005) Loss- and gain-of-function analysis of the lipid raft proteins Reggie/Flotillin in Drosophila: they are

posttranslationally regulated, and misexpression interferes with wing and eye development. *Mol Cell Neurosci* 30(3): 326-338

Honda R, Korner R, Nigg EA (2003) Exploring the functional interactions between Aurora B, INCENP, and survivin in mitosis. *Mol Biol Cell* 14(8): 3325-3341

Hooper C, Chapple JP, Lovestone S, Killick R (2007) The Notch-1 intracellular domain is found in sub-nuclear bodies in SH-SY5Y neuroblastomas and in primary cortical neurons. *Neurosci Lett* 415(2): 135-139

Hsieh JJ, Zhou S, Chen L, Young DB, Hayward SD (1999) CIR, a corepressor linking the DNA binding factor CBF1 to the histone deacetylase complex. *Proc Natl Acad Sci U S A* 96(1): 23-28

Hu W, Kemp BE, Jans DA (2005) Kinetic properties of nuclear transport conferred by the retinoblastoma (Rb) NLS. *J Cell Biochem* 95(4): 782-793

Huang CY, Lu TY, Bair CH, Chang YS, Jwo JK, Chang W (2008) A novel cellular protein, VPEF, facilitates vaccinia virus penetration into HeLa cells through fluid phase endocytosis. *J Virol* 82(16): 7988-7999

Hung AY, Sheng M (2002) PDZ domains: structural modules for protein complex assembly. *J Biol Chem* 277(8): 5699-5702

I

Ikui AE, Furuya K, Yanagida M, Matsumoto T (2002) Control of localization of a spindle checkpoint protein, Mad2, in fission yeast. *J Cell Sci* 115(Pt 8): 1603-1610

Incardona JP, Eaton S (2000) Cholesterol in signal transduction. *Curr Opin Cell Biol* 12(2): 193-203

Iso T, Kedes L, Hamamori Y (2003) HES and HERP families: multiple effectors of the Notch signaling pathway. *J Cell Physiol* 194(3): 237-255

J

Janes PW, Ley SC, Magee AI, Kabouridis PS (2000) The role of lipid rafts in T cell antigen receptor (TCR) signalling. *Semin Immunol* 12(1): 23-34

Jeyaprakash AA, Klein UR, Lindner D, Ebert J, Nigg EA, Conti E (2007) Structure of a Survivin-Borealin-INCENP core complex reveals how chromosomal passengers travel together. *Cell* 131(2): 271-285

Joshi B, Rastogi S, Morris M, Carastro LM, DeCook C, Seto E, Chellappan SP (2007) Differential regulation of human YY1 and caspase 7 promoters by prohibitin through E2F1 and p53 binding sites. *Biochem J* 401(1): 155-166

Juan G, Traganos F, James WM, Ray JM, Roberge M, Sauve DM, Anderson H, Darzynkiewicz Z (1998) Histone H3 phosphorylation and expression of cyclins A and B1 measured in individual cells during their progression through G2 and mitosis. *Cytometry* 32(2): 71-77

K

Kadesch T (2004) Notch signaling: the demise of elegant simplicity. *Curr Opin Genet Dev* 14(5): 506-512

Kamakura S, Oishi K, Yoshimatsu T, Nakafuku M, Masuyama N, Gotoh Y (2004) Hes binding to STAT3 mediates crosstalk between Notch and JAK-STAT signalling. *Nat Cell Biol* 6(6): 547-554

Kao HY, Ordentlich P, Koyano-Nakagawa N, Tang Z, Downes M, Kintner CR, Evans RM, Kadesch T (1998) A histone deacetylase corepressor complex regulates the Notch signal transduction pathway. *Genes Dev* 12(15): 2269-2277

Katic I, Vallier LG, Greenwald I (2005) New positive regulators of lin-12 activity in *Caenorhabditis elegans* include the BRE-5/Brainiac glycosphingolipid biosynthesis enzyme. *Genetics* 171(4): 1605-1615

Kato N, Nakanishi M, Hirashima N (2006) Flotillin-1 regulates IgE receptor-mediated signaling in rat basophilic leukemia (RBL-2H3) cells. *J Immunol* 177(1): 147-154

Kelly AE, Sampath SC, Maniar TA, Woo EM, Chait BT, Funabiki H (2007) Chromosomal enrichment and activation of the Aurora B pathway are coupled to spatially regulate spindle assembly. *Dev Cell* 12(1): 31-43

Kim KB, Crews CM (2008) Chemical genetics: exploring the role of the proteasome in cell biology using natural products and other small molecule proteasome inhibitors. *J Med Chem* 51(9): 2600-2605

Kim SH, Ikeuchi T, Yu C, Sisodia SS (2003) Regulated hyperaccumulation of presenilin-1 and the "gamma-secretase" complex. Evidence for differential intramembranous processing of transmembrane substrates. *J Biol Chem* 278(36): 33992-34002

Kimura A, Baumann CA, Chiang SH, Saltiel AR (2001) The sorbin homology domain: a motif for the targeting of proteins to lipid rafts. *Proc Natl Acad Sci U S A* 98(16): 9098-9103

Kirchhausen T (1999) Adaptors for clathrin-mediated traffic. *Annu Rev Cell Dev Biol* 15: 705-732

Kopan R, Nye JS, Weintraub H (1994) The intracellular domain of mouse Notch: a constitutively activated repressor of myogenesis directed at the basic helix-loop-helix region of MyoD. *Development* 120(9): 2385-2396

Kotani S, Tugendreich S, Fujii M, Jorgensen PM, Watanabe N, Hoog C, Hieter P, Todokoro K (1998) PKA and MPF-activated polo-like kinase regulate anaphase-promoting complex activity and mitosis progression. *Mol Cell* 1(3): 371-380

Kovall RA (2007) Structures of CSL, Notch and Mastermind proteins: piecing together an active transcription complex. *Curr Opin Struct Biol* 17(1): 117-127

Kramer ER, Scheuringer N, Podtelejnikov AV, Mann M, Peters JM (2000) Mitotic regulation of the APC activator proteins CDC20 and CDH1. *Mol Biol Cell* 11(5): 1555-1569

Kunitoku N, Sasayama T, Marumoto T, Zhang D, Honda S, Kobayashi O, Hatakeyama K, Ushio Y, Saya H, Hirota T (2003) CENP-A phosphorylation by Aurora-A in prophase is required for enrichment of Aurora-B at inner centromeres and for kinetochore function. *Dev Cell* 5(6): 853-864

L

Lahav-Baratz S, Sudakin V, Ruderman JV, Hershko A (1995) Reversible phosphorylation controls the activity of cyclosome-associated cyclin-ubiquitin ligase. *Proc Natl Acad Sci U S A* 92(20): 9303-9307

Lai EC (2002) Keeping a good pathway down: transcriptional repression of Notch pathway target genes by CSL proteins. *EMBO Rep* 3(9): 840-845

Lamaze C, Dujeancourt A, Baba T, Lo CG, Benmerah A, Dautry-Varsat A (2001) Interleukin 2 receptors and detergent-resistant membrane domains define a clathrin-independent endocytic pathway. *Mol Cell* 7(3): 661-671

Lampson MA, Kapoor TM (2005) The human mitotic checkpoint protein BubR1 regulates chromosome-spindle attachments. *Nat Cell Biol* 7(1): 93-98

Lang DM, Lommel S, Jung M, Ankerhold R, Petrausch B, Laessing U, Wiechers MF, Plattner H, Stuermer CA (1998) Identification of Reggie-1 and Reggie-2 as plasmamembrane-associated proteins which cocluster with activated GPI-anchored cell adhesion molecules in non-caveolar micropatches in neurons. *J Neurobiol* 37(4): 502-523

Lange BM, Rebollo E, Herold A, Gonzalez C (2002) Cdc37 is essential for chromosome segregation and cytokinesis in higher eukaryotes. *EMBO J* 21(20): 5364-5374

Langhorst MF, Reuter A, Stuermer CA (2005) Scaffolding microdomains and beyond: the function of Reggie/Flotillin proteins. *Cell Mol Life Sci* 62(19-20): 2228-2240

Le Borgne R, Bardin A, Schweigert F (2005) The roles of receptor and ligand endocytosis in regulating Notch signaling. *Development* 132(8): 1751-1762

Legler DF, Micheau O, Doucey MA, Tschopp J, Bron C (2003) Recruitment of TNF receptor 1 to lipid rafts is essential for TNFalpha-mediated NF-kappaB activation. *Immunity* 18(5): 655-664

Lens SM, Wolthuis RM, Klompmaker R, Kauw J, Agami R, Brummelkamp T, Kops G, Medema RH (2003) Survivin is required for a sustained spindle checkpoint arrest in response to lack of tension. *EMBO J* 22(12): 2934-2947

Liu J, Deyoung SM, Zhang M, Dold LH, Saltiel AR (2005) The Stomatin/prohibitin/Flotillin/HflK/C domain of Flotillin-1 contains distinct sequences that direct plasma membrane localization and protein interactions in 3T3-L1 adipocytes. *J Biol Chem* 280(16): 16125-16134

Lodish HF, Berk A, Lawrence Zipurski S, Matsudaira P, Baltimore D, Darnell J (1999) *Molecular Cell Biology*.

Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193(1): 265-275

Lu H, Bilder D (2005) Endocytic control of epithelial polarity and proliferation in Drosophila. *Nat Cell Biol* 7(12): 1232-1239

M

Ma Y, Chan CY, He ML (2007) RNA interference and antiviral therapy. *World J Gastroenterol* 13(39): 5169-5179

Marquez DC, Chen HW, Curran EM, Welshons WV, Pietras RJ (2006) Estrogen receptors in membrane lipid rafts and signal transduction in breast cancer. *Mol Cell Endocrinol* 246(1-2): 91-100

Martinez Arias A, Zecchini V, Brennan K (2002) CSL-independent Notch signalling: a checkpoint in cell fate decisions during development? *Curr Opin Genet Dev* 12(5): 524-533

Mastick CC, Brady MJ, Saltiel AR (1995) Insulin stimulates the tyrosine phosphorylation of Caveolin. *J Cell Biol* 129(6): 1523-1531

Matsuno K, Diederich RJ, Go MJ, Blaumueller CM, Artavanis-Tsakonas S (1995) Deltex acts as a positive regulator of Notch signaling through interactions with the Notch ankyrin repeats. *Development* 121(8): 2633-2644

Meyer O (2002) [CREST syndrome]. *Ann Med Interne (Paris)* 153(3): 183-188

Millband DN, Campbell L, Hardwick KG (2002) The awesome power of multiple model systems: interpreting the complex nature of spindle checkpoint signaling. *Trends Cell Biol* 12(5): 205-209

Minshall RD, Tiruppathi C, Vogel SM, Niles WD, Gilchrist A, Hamm HE, Malik AB (2000) Endothelial cell-surface gp60 activates vesicle formation and trafficking via G(i)-coupled Src kinase signaling pathway. *J Cell Biol* 150(5): 1057-1070

Mittler G, Stuhler T, Santolin L, Uhlmann T, Kremmer E, Lottspeich F, Berti L, Meisterernst M (2003) A novel docking site on Mediator is critical for activation by VP16 in mammalian cells. *EMBO J* 22(24): 6494-6504

Mohr OL (1919) Character Changes Caused by Mutation of an Entire Region of a Chromosome in Drosophila. *Genetics* 4(3): 275-282

Morais VA, Leight S, Pijak DS, Lee VM, Costa J (2008) Cellular localization of Nicastrin affects amyloid beta species production. *FEBS Lett* 582(3): 427-433

Morgan DO (1997) Cyclin-dependent kinases: engines, clocks, and microprocessors. *Annu Rev Cell Dev Biol* 13: 261-291

Morgan JR, Augustine GJ, Lafer EM (2002) Synaptic vesicle endocytosis: the races, places, and molecular faces. *Neuromolecular Med* 2(2): 101-114

Morrison C, Henzing AJ, Jensen ON, Osheroff N, Dodson H, Kandels-Lewis SE, Adams RR, Earnshaw WC (2002) Proteomic analysis of human metaphase chromosomes reveals topoisomerase II alpha as an Aurora B substrate. *Nucleic Acids Res* 30(23): 5318-5327

Morrow IC, Rea S, Martin S, Prior IA, Prohaska R, Hancock JF, James DE, Parton RG (2002) Flotillin-1/Reggie-2 traffics to surface raft domains via a novel golgi-independent pathway. Identification of a novel membrane targeting domain and a role for palmitoylation. *J Biol Chem* 277(50): 48834-48841

Mukhopadhyay D, Riezman H (2007) Proteasome-independent functions of ubiquitin in endocytosis and signaling. *Science* 315(5809): 201-205

Mumm JS, Kopan R (2000) Notch signaling: from the outside in. *Dev Biol* 228(2): 151-165

Mumm JS, Schroeter EH, Saxena MT, Griesemer A, Tian X, Pan DJ, Ray WJ, Kopan R (2000) A ligand-induced extracellular cleavage regulates gamma-secretase-like proteolytic activation of Notch1. *Mol Cell* 5(2): 197-206

N

Nabi IR, Le PU (2003) Caveolae/raft-dependent endocytosis. *J Cell Biol* 161(4): 673-677

Nasmyth K (1999) Separating sister chromatids. *Trends Biochem Sci* 24(3): 98-104

Neumann-Giesen C, Falkenbach B, Beicht P, Claassen S, Luers G, Stuermer CA, Herzog V, Tikkanen R (2004) Membrane and raft association of Reggie-1/Flotillin-2: role of

myristoylation, palmitoylation and oligomerization and induction of filopodia by overexpression. *Biochem J* 378(Pt 2): 509-518

Nguyen HG, Chinnappan D, Urano T, Ravid K (2005) Mechanism of Aurora-B degradation and its dependency on intact KEN and A-boxes: identification of an aneuploidy-promoting property. *Mol Cell Biol* 25(12): 4977-4992

Nichols B (2003) Caveosomes and endocytosis of lipid rafts. *J Cell Sci* 116(Pt 23): 4707-4714

Nichols BJ, Lippincott-Schwartz J (2001) Endocytosis without clathrin coats. *Trends Cell Biol* 11(10): 406-412

Nichols JT, Miyamoto A, Weinmaster G (2007) Notch signaling--constantly on the move. *Traffic* 8(8): 959-969

Nijtmans LG, de Jong L, Artal Sanz M, Coates PJ, Berden JA, Back JW, Muijsers AO, van der Spek H, Grivell LA (2000) Prohibitins act as a membrane-bound chaperone for the stabilization of mitochondrial proteins. *EMBO J* 19(11): 2444-2451

Nizzari M, Venezia V, Bianchini P, Caorsi V, Diaspro A, Repetto E, Thellung S, Corsaro A, Carlo P, Schettini G, Florio T, Russo C (2007) Amyloid precursor protein and Presenilin 1 interaction studied by FRET in human H4 cells. *Ann N Y Acad Sci* 1096: 249-257

Nye JS, Kopan R, Axel R (1994) An activated Notch suppresses neurogenesis and myogenesis but not gliogenesis in mammalian cells. *Development* 120(9): 2421-2430

O

Oh P, McIntosh DP, Schnitzer JE (1998) Dynamin at the neck of caveolae mediates their budding to form transport vesicles by GTP-driven fission from the plasma membrane of endothelium. *J Cell Biol* 141(1): 101-114

Okajima T, Matsuura A, Matsuda T (2008) Biological functions of glycosyltransferase genes involved in O-fucose glycan synthesis. *J Biochem* 144(1): 1-6

Okamoto T, Schlegel A, Scherer PE, Lisanti MP (1998) Caveolins, a family of scaffolding proteins for organizing "preassembled signaling complexes" at the plasma membrane. *J Biol Chem* 273(10): 5419-5422

Ota T, Suto S, Katayama H, Han ZB, Suzuki F, Maeda M, Tanino M, Terada Y, Tatsuka M (2002) Increased mitotic phosphorylation of histone H3 attributable to AIM-1/Aurora-B overexpression contributes to chromosome number instability. *Cancer Res* 62(18): 5168-5177

P

Palomero T, Sulis ML, Cortina M, Real PJ, Barnes K, Ciofani M, Caparros E, Buteau J, Brown K, Perkins SL, Bhagat G, Agarwal AM, Basso G, Castillo M, Nagase S, Cordon-Cardo C, Parsons R, Zuniga-Pflucker JC, Dominguez M, Ferrando AA (2007) Mutational loss of PTEN induces resistance to NOTCH1 inhibition in T-cell leukemia. *Nat Med* 13(10): 1203-1210

Parton RG, Hanzal-Bayer M, Hancock JF (2006) Biogenesis of caveolae: a structural model for Caveolin-induced domain formation. *J Cell Sci* 119(Pt 5): 787-796

Parton RG, Simons K (2007) The multiple faces of caveolae. *Nat Rev Mol Cell Biol* 8(3): 185-194

Pasternak SH, Bagshaw RD, Guiral M, Zhang S, Ackerley CA, Pak BJ, Callahan JW, Mahuran DJ (2003) Presenilin-1, nicastrin, amyloid precursor protein, and gamma-secretase activity are co-localized in the lysosomal membrane. *J Biol Chem* 278(29): 26687-26694

Payne CK, Jones SA, Chen C, Zhuang X (2007) Internalization and trafficking of cell surface proteoglycans and proteoglycan-binding ligands. *Traffic* 8(4): 389-401

Pelkmans L, Burli T, Zerial M, Helenius A (2004) Caveolin-stabilized membrane domains as multifunctional transport and sorting devices in endocytic membrane traffic. *Cell* 118(6): 767-780

Pelkmans L, Kartenbeck J, Helenius A (2001) Caveolar endocytosis of simian virus 40 reveals a new two-step vesicular-transport pathway to the ER. *Nat Cell Biol* 3(5): 473-483

Pelkmans L, Puntener D, Helenius A (2002) Local actin polymerization and dynamin recruitment in SV40-induced internalization of caveolae. *Science* 296(5567): 535-539

Pierchala BA, Milbrandt J, Johnson EM, Jr. (2006) Glial cell line-derived neurotrophic factor-dependent recruitment of Ret into lipid rafts enhances signaling by partitioning Ret from proteasome-dependent degradation. *J Neurosci* 26(10): 2777-2787

Poulson DF (1937) Chromosomal Deficiencies and the Embryonic Development of *Drosophila Melanogaster*. *Proc Natl Acad Sci U S A* 23(3): 133-137

Pralle A, Keller P, Florin EL, Simons K, Horber JK (2000) Sphingolipid-cholesterol rafts diffuse as small entities in the plasma membrane of mammalian cells. *J Cell Biol* 148(5): 997-1008

Prigent C, Dimitrov S (2003) Phosphorylation of serine 10 in histone H3, what for? *J Cell Sci* 116(Pt 18): 3677-3685

R

Rajendran L, Masilamani M, Solomon S, Tikkanen R, Stuermer CA, Plattner H, Illges H (2003) Asymmetric localization of Flotillins/Reggies in preassembled platforms confers inherent polarity to hematopoietic cells. *Proc Natl Acad Sci U S A* 100(14): 8241-8246

Rajendran L, Simons K (2005) Lipid rafts and membrane dynamics. *J Cell Sci* 118(Pt 6): 1099-1102

Rand MD, Grimm LM, Artavanis-Tsakonas S, Patriub V, Blacklow SC, Sklar J, Aster JC (2000) Calcium depletion dissociates and activates heterodimeric notch receptors. *Mol Cell Biol* 20(5): 1825-1835

Rastogi S, Joshi B, Dasgupta P, Morris M, Wright K, Chellappan S (2006) Prohibitin facilitates cellular senescence by recruiting specific corepressors to inhibit E2F target genes. *Mol Cell Biol* 26(11): 4161-4171

Ray WJ, Yao M, Mumm J, Schroeter EH, Saftig P, Wolfe M, Selkoe DJ, Kopan R, Goate AM (1999) Cell surface presenilin-1 participates in the gamma-secretase-like proteolysis of Notch. *J Biol Chem* 274(51): 36801-36807

Resh MD (1999) Fatty acylation of proteins: new insights into membrane targeting of myristoylated and palmitoylated proteins. *Biochim Biophys Acta* 1451(1): 1-16

Rivera-Milla E, Stuermer CA, Malaga-Trillo E (2006) Ancient origin of Reggie (Flotillin), Reggie-like, and other lipid-raft proteins: convergent evolution of the SPFH domain. *Cell Mol Life Sci* 63(3): 343-357

Ronchini C, Capobianco AJ (2001) Induction of cyclin D1 transcription and CDK2 activity by Notch(ic): implication for cell cycle disruption in transformation by Notch(ic). *Mol Cell Biol* 21(17): 5925-5934

Rothberg KG, Heuser JE, Donzell WC, Ying YS, Glenney JR, Anderson RG (1992) Caveolin, a protein component of caveolae membrane coats. *Cell* 68(4): 673-682

Roy S, Luetterforst R, Harding A, Apolloni A, Etheridge M, Stang E, Rolls B, Hancock JF, Parton RG (1999) Dominant-negative Caveolin inhibits H-Ras function by disrupting cholesterol-rich plasma membrane domains. *Nat Cell Biol* 1(2): 98-105

Ruchaud S, Carmena M, Earnshaw WC (2007) Chromosomal passengers: conducting cell division. *Nat Rev Mol Cell Biol* 8(10): 798-812

S

Sabharanjak S, Sharma P, Parton RG, Mayor S (2002) GPI-anchored proteins are delivered to recycling endosomes via a distinct cdc42-regulated, clathrin-independent pinocytic pathway. *Dev Cell* 2(4): 411-423

- Salah SM, Nasmyth K (2000) Destruction of the securin Pds1p occurs at the onset of anaphase during both meiotic divisions in yeast. *Chromosoma* 109(1-2): 27-34
- Salanueva IJ, Cerezo A, Guadamilas MC, del Pozo MA (2007) Integrin regulation of Caveolin function. *J Cell Mol Med* 11(5): 969-980
- Sanchez-Irizarry C, Carpenter AC, Weng AP, Pear WS, Aster JC, Blacklow SC (2004) Notch subunit heterodimerization and prevention of ligand-independent proteolytic activation depend, respectively, on a novel domain and the LNR repeats. *Mol Cell Biol* 24(21): 9265-9273
- Santamaria A, Castellanos E, Gomez V, Benedit P, Renau-Piqueras J, Morote J, Reventos J, Thomson TM, Paciucci R (2005) PTOV1 enables the nuclear translocation and mitogenic activity of Flotillin-1, a major protein of lipid rafts. *Mol Cell Biol* 25(5): 1900-1911
- Sasaki Y, Oshima Y, Koyama R, Maruyama R, Akashi H, Mita H, Toyota M, Shinomura Y, Imai K, Tokino T (2008) Identification of Flotillin-2, a major protein on lipid rafts, as a novel target of p53 family members. *Mol Cancer Res* 6(3): 395-406
- Scheiffele P, Peranen J, Simons K (1995) N-glycans as apical sorting signals in epithelial cells. *Nature* 378(6552): 96-98
- Schneider A, Rajendran L, Honsho M, Gralle M, Donnert G, Wouters F, Hell SW, Simons M (2008) Flotillin-dependent clustering of the amyloid precursor protein regulates its endocytosis and amyloidogenic processing in neurons. *J Neurosci* 28(11): 2874-2882
- Scholzen T, Gerdes J (2000) The Ki-67 protein: from the known and the unknown. *J Cell Physiol* 182(3): 311-322
- Schroeter EH, Kisslinger JA, Kopan R (1998) Notch-1 signalling requires ligand-induced proteolytic release of intracellular domain. *Nature* 393(6683): 382-386
- Schuck S, Simons K (2004) Polarized sorting in epithelial cells: raft clustering and the biogenesis of the apical membrane. *J Cell Sci* 117(Pt 25): 5955-5964
- Schulte T, Paschke KA, Laessing U, Lottspeich F, Stuermer CA (1997) Reggie-1 and Reggie-2, two cell surface proteins expressed by retinal ganglion cells during axon regeneration. *Development* 124(2): 577-587
- Schwab M, Lutum AS, Seufert W (1997) Yeast Hct1 is a regulator of Clb2 cyclin proteolysis. *Cell* 90(4): 683-693
- Seugnet L, Simpson P, Haenlin M (1997) Requirement for dynamin during Notch signaling in Drosophila neurogenesis. *Dev Biol* 192(2): 585-598

- Shawber C, Nofziger D, Hsieh JJ, Lindsell C, Bogler O, Hayward D, Weinmaster G (1996) Notch signaling inhibits muscle cell differentiation through a CBF1-independent pathway. *Development* 122(12): 3765-3773
- Sheets ED, Holowka D, Baird B (1999) Membrane organization in immunoglobulin E receptor signaling. *Curr Opin Chem Biol* 3(1): 95-99
- Shirayama M, Toth A, Galova M, Nasmyth K (1999) APC(Cdc20) promotes exit from mitosis by destroying the anaphase inhibitor Pds1 and cyclin Clb5. *Nature* 402(6758): 203-207
- Simons K, Ikonen E (1997) Functional rafts in cell membranes. *Nature* 387(6633): 569-572
- Simons K, Toomre D (2000) Lipid rafts and signal transduction. *Nat Rev Mol Cell Biol* 1(1): 31-39
- Sisodia SS, Annaert W, Kim SH, De Strooper B (2001) Gamma-secretase: never more enigmatic. *Trends Neurosci* 24(11 Suppl): S2-6
- Slaughter N, Laux I, Tu X, Whitelegge J, Zhu X, Effros R, Bickel P, Nel A (2003) The Flotillins are integral membrane proteins in lipid rafts that contain TCR-associated signaling components: implications for T-cell activation. *Clin Immunol* 108(2): 138-151
- Snyders L, Umlauf E, Prohaska R (1998) Oligomeric nature of the integral membrane protein Stomatin. *J Biol Chem* 273(27): 17221-17226
- Solis GP, Hoegg M, Munderloh C, Schrock Y, Malaga-Trillo E, Rivera-Milla E, Stuermer CA (2007) Reggie/Flotillin proteins are organized into stable tetramers in membrane microdomains. *Biochem J* 403(2): 313-322
- Song J, Salek-Ardakani S, So T, Croft M (2007) The kinases Aurora B and mTOR regulate the G1-S cell cycle progression of T lymphocytes. *Nat Immunol* 8(1): 64-73
- Soond SM, Everson B, Riches DW, Murphy G (2005) ERK-mediated phosphorylation of Thr735 in TNFalpha-converting enzyme and its potential role in TACE protein trafficking. *J Cell Sci* 118(Pt 11): 2371-2380
- Steglich G, Neupert W, Langer T (1999) Prohibitins regulate membrane protein degradation by the m-AAA protease in mitochondria. *Mol Cell Biol* 19(5): 3435-3442
- Stewart S, Fang G (2005) Destruction box-dependent degradation of Aurora B is mediated by the anaphase-promoting complex/cyclosome and Cdh1. *Cancer Res* 65(19): 8730-8735
- Stuermer CA, Lang DM, Kirsch F, Wiechers M, Deininger SO, Plattner H (2001) Glycosylphosphatidyl inositol-anchored proteins and fyn kinase assemble in noncaveolar plasma membrane microdomains defined by Reggie-1 and -2. *Mol Biol Cell* 12(10): 3031-3045

Stuermer CA, Langhorst MF, Wiechers MF, Legler DF, Von Hanwehr SH, Guse AH, Plattner H (2004) PrPc capping in T cells promotes its association with the lipid raft proteins Reggie-1 and Reggie-2 and leads to signal transduction. *FASEB J* 18(14): 1731-1733

Sudakin V, Chan GK, Yen TJ (2001) Checkpoint inhibition of the APC/C in HeLa cells is mediated by a complex of BUBR1, BUB3, CDC20, and MAD2. *J Cell Biol* 154(5): 925-936

Sugawara Y, Nishii H, Takahashi T, Yamauchi J, Mizuno N, Tago K, Itoh H (2007) The lipid raft proteins Flotillins/Reggies interact with Galphaq and are involved in Gq-mediated p38 mitogen-activated protein kinase activation through tyrosine kinase. *Cell Signal* 19(6): 1301-1308

Sumara I, Peter M (2007) A Cul3-based E3 ligase regulates mitosis and is required to maintain the spindle assembly checkpoint in human cells. *Cell Cycle* 6(24): 3004-3010

Sumara I, Vorlaufer E, Stukenberg PT, Kelm O, Redemann N, Nigg EA, Peters JM (2002) The dissociation of cohesin from chromosomes in prophase is regulated by Polo-like kinase. *Mol Cell* 9(3): 515-525

Sun L, Gao J, Dong X, Liu M, Li D, Shi X, Dong JT, Lu X, Liu C, Zhou J (2008) EB1 promotes Aurora-B kinase activity through blocking its inactivation by protein phosphatase 2A. *Proc Natl Acad Sci U S A* 105(20): 7153-7158

Sundaram MV (2005) The love-hate relationship between Ras and Notch. *Genes Dev* 19(16): 1825-1839

T

Tagawa A, Mezzacasa A, Hayer A, Longatti A, Pelkmans L, Helenius A (2005) Assembly and trafficking of caveolar domains in the cell: caveolae as stable, cargo-triggered, vesicular transporters. *J Cell Biol* 170(5): 769-779

Tanaka TU, Rachidi N, Janke C, Pereira G, Galova M, Schiebel E, Stark MJ, Nasmyth K (2002) Evidence that the Ipl1-Sli15 (Aurora kinase-INCENP) complex promotes chromosome bi-orientation by altering kinetochore-spindle pole connections. *Cell* 108(3): 317-329

Tang Z, Bharadwaj R, Li B, Yu H (2001) Mad2-Independent inhibition of APCCdc20 by the mitotic checkpoint protein BubR1. *Dev Cell* 1(2): 227-237

Tang Z, Scherer PE, Okamoto T, Song K, Chu C, Kohtz DS, Nishimoto I, Lodish HF, Lisanti MP (1996) Molecular cloning of Caveolin-3, a novel member of the Caveolin gene family expressed predominantly in muscle. *J Biol Chem* 271(4): 2255-2261

Tansey MG, Baloh RH, Milbrandt J, Johnson EM, Jr. (2000) GFRalpha-mediated localization of RET to lipid rafts is required for effective downstream signaling, differentiation, and neuronal survival. *Neuron* 25(3): 611-623

Tavernarakis N, Driscoll M, Kyrides NC (1999) The SPFH domain: implicated in regulating targeted protein turnover in Stomatins and other membrane-associated proteins. *Trends Biochem Sci* 24(11): 425-427

Taylor WR, Stark GR (2001) Regulation of the G2/M transition by p53. *Oncogene* 20(15): 1803-1815

Terada Y, Tatsuka M, Suzuki F, Yasuda Y, Fujita S, Otsu M (1998) AIM-1: a mammalian midbody-associated protein required for cytokinesis. *EMBO J* 17(3): 667-676

Thompson HM, Skop AR, Euteneuer U, Meyer BJ, McNiven MA (2002) The large GTPase dynamin associates with the spindle midzone and is required for cytokinesis. *Curr Biol* 12(24): 2111-2117

Towbin H, Staehelin T, Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci U S A* 76(9): 4350-4354

Tulu US, Fagerstrom C, Ferenz NP, Wadsworth P (2006) Molecular requirements for kinetochore-associated microtubule formation in mammalian cells. *Curr Biol* 16(5): 536-541

U

Urano Y, Hayashi I, Isso N, Reid PC, Shibasaki Y, Noguchi N, Tomita T, Iwatsubo T, Hamakubo T, Kodama T (2005) Association of active gamma-secretase complex with lipid rafts. *J Lipid Res* 46(5): 904-912

V

Vaccari T, Lu H, Kanwar R, Fortini ME, Bilder D (2008) Endosomal entry regulates Notch receptor activation in *Drosophila melanogaster*. *J Cell Biol* 180(4): 755-762

Vader G, Medema RH, Lens SM (2006) The chromosomal passenger complex: guiding Aurora-B through mitosis. *J Cell Biol* 173(6): 833-837

Van Damme D, Inze D, Russinova E (2008) Vesicle trafficking during somatic cytokinesis. *Plant Physiol* 147(4): 1544-1552

Vetrivel KS, Cheng H, Kim SH, Chen Y, Barnes NY, Parent AT, Sisodia SS, Thinakaran G (2005) Spatial segregation of gamma-secretase and substrates in distinct membrane domains. *J Biol Chem* 280(27): 25892-25900

Visintin R, Prinz S, Amon A (1997) CDC20 and CDH1: a family of substrate-specific activators of APC-dependent proteolysis. *Science* 278(5337): 460-463

Voglmaier SM, Kam K, Yang H, Fortin DL, Hua Z, Nicoll RA, Edwards RH (2006) Distinct endocytic pathways control the rate and extent of synaptic vesicle protein recycling. *Neuron* 51(1): 71-84

Volonte D, Galbiati F, Li S, Nishiyama K, Okamoto T, Lisanti MP (1999) Flotillins/cavatellins are differentially expressed in cells and tissues and form a hetero-oligomeric complex with Caveolins in vivo. Characterization and epitope-mapping of a novel Flotillin-1 monoclonal antibody probe. *J Biol Chem* 274(18): 12702-12709

Vong QP, Cao K, Li HY, Iglesias PA, Zheng Y (2005) Chromosome alignment and segregation regulated by ubiquitination of survivin. *Science* 310(5753): 1499-1504

W

Wadsworth P, Khodjakov A (2004) E pluribus unum: towards a universal mechanism for spindle assembly. *Trends Cell Biol* 14(8): 413-419

Wang S, Fusaro G, Padmanabhan J, Chellappan SP (2002) Prohibitin co-localizes with Rb in the nucleus and recruits N-CoR and HDAC1 for transcriptional repression. *Oncogene* 21(55): 8388-8396

Wassmann K, Benezra R (1998) Mad2 transiently associates with an APC/p55Cdc complex during mitosis. *Proc Natl Acad Sci U S A* 95(19): 11193-11198

Weijzen S, Rizzo P, Braid M, Vaishnav R, Jonkheer SM, Zlobin A, Osborne BA, Gottipati S, Aster JC, Hahn WC, Rudolf M, Siziopikou K, Kast WM, Miele L (2002) Activation of Notch-1 signaling maintains the neoplastic phenotype in human Ras-transformed cells. *Nat Med* 8(9): 979-986

Weinmaster G, Roberts VJ, Lemke G (1992) Notch2: a second mammalian Notch gene. *Development* 116(4): 931-941

Wharton KA, Johansen KM, Xu T, Artavanis-Tsakonas S (1985) Nucleotide sequence from the neurogenic locus notch implies a gene product that shares homology with proteins containing EGF-like repeats. *Cell* 43(3 Pt 2): 567-581

Wheatley SP, Carvalho A, Vagnarelli P, Earnshaw WC (2001) INCENP is required for proper targeting of Survivin to the centromeres and the anaphase spindle during mitosis. *Curr Biol* 11(11): 886-890

Wood DR, Nye JS, Lamb NJ, Fernandez A, Kitzmann M (2005) Intracellular retention of Caveolin 1 in presenilin-deficient cells. *J Biol Chem* 280(8): 6663-6668

Wu L, Sun T, Kobayashi K, Gao P, Griffin JD (2002) Identification of a family of mastermind-like transcriptional coactivators for mammalian notch receptors. *Mol Cell Biol* 22(21): 7688-7700

Y

Yamamoto M, Toya Y, Schwencke C, Lisanti MP, Myers MG, Jr., Ishikawa Y (1998) Caveolin is an activator of insulin receptor signaling. *J Biol Chem* 273(41): 26962-26968

Yamamoto TM, Lewellyn AL, Maller JL (2008) Regulation of the Aurora B chromosome passenger protein complex during oocyte maturation in *Xenopus laevis*. *Mol Cell Biol* 28(12): 4196-4203

Z

Zhang J, Ye W, Wang R, Wolfe MS, Greenberg BD, Selkoe DJ (2002) Proteolysis of chimeric beta-amyloid precursor proteins containing the Notch transmembrane domain yields amyloid beta-like peptides. *J Biol Chem* 277(17): 15069-15075

Zheng N, Schulman BA, Song L, Miller JJ, Jeffrey PD, Wang P, Chu C, Koepp DM, Elledge SJ, Pagano M, Conaway RC, Conaway JW, Harper JW, Pavletich NP (2002) Structure of the Cul1-Rbx1-Skp1-F boxSkp2 SCF ubiquitin ligase complex. *Nature* 416(6882): 703-709

Zhou BB, Elledge SJ (2000) The DNA damage response: putting checkpoints in perspective. *Nature* 408(6811): 433-439

Zhou S, Fujimuro M, Hsieh JJ, Chen L, Miyamoto A, Weinmaster G, Hayward SD (2000) SKIP, a CBF1-associated protein, interacts with the ankyrin repeat domain of NotchIC To facilitate NotchIC function. *Mol Cell Biol* 20(7): 2400-2410