

VI.BIBLIOGRAFIA

- Abdulaev N. G. (2003) Building a stage for interhelical play in rhodopsin. *Trends Biochem. Sci.* **28**; 399-402.
- Alberts B., Bray P., Lewis J., Raff M., Roberts R. i Watson J. D. *Molecular Biology of the cell* (1994). 3th edition. Garland Publishing, Inc.
- Alewijnse A. E., Timmerman H., Jacobs E. H., Smit M. J., Roovers E., Cotecchia S. i Leurs R. (2000) The effect of mutations in the DRY motif on the constitutive activity and structural instability of the histamine H(2) receptor. *Mol. Pharmacol.* **57**; 890-898.
- al-Jandal N., Farrar G. J., Kiang A. S., Humphries M. M., Bannon N., Findlay J. B., Humphries P. i Kenna P. F. (1999) A novel mutation within the rhodopsin gene (Thr-94-Ile) causing autosomal dominant congenital stationary night blindness. *Hum. Mutat.* **13**; 75-81.
- Andres A., Garriga P. i Manyosa J. (2003) Altered functionality in rhodopsin point mutants associated with retinitis pigmentosa. *Biochem. Biophys. Res. Commun.* **303**; 294-301.
- Anzenbacherova E., Bec N., Anzenbacher P., Hudecek J., Soucek P., Jung C., Munro A. W. i Lange R. (2000) Flexibility and stability of the structure of cytochromes P450 3A4 and BM-3. *Eur. J. Biochem.* **267**; 2916-2920.
- Arnis S., Fahmy K., Hofmann K. P. i Sakmar, T. P. (1994) A conserved carboxylic acid group mediates light-dependent proton uptake and signaling by rhodopsin. *J. Biol. Chem.* **269**; 23879-23881.
- Arora K. K., Cheng Z. i Catt K. J. (1997) Mutations of the conserved DRS motif in the second intracellular loop of the gonadotropin-releasing hormone receptor affect expression, activation, and internalization. *Mol. Endocrinol.* **11**; 203-212.
- Ashwood-Smith M. J. (1967) Current concept concerning radioprotective and cryoprotective properties of dimethyl sulfoxide in cellular systems. *Ann. N. Y. Acad. Sci. USA* **141**; 41-62.
- Atwood C. S., Moir R. D. Huang X., Scarpa R. C., Bacarra N. M., Romano D. M., Hartshorn M. A., Tanzi R. E. i Bush A. I. (1998) Dramatic aggregation of Alzheimer A β by Cu(II) is induced by conditions representing physiological acidosis. *J. Biol. Chem.* **273**; 12817-18226.
- Ballesteros J. A. i Weinstein H. (1995) Integrated Methods for the Construction of Three-Dimensional Models and Computational Probing of Structure-Function Relations in G Protein-Coupled Receptors. *Methods Neuroscience* **25**; 366-428.
- Ballesteros J. A., Kitanovic S., Guarnieri F., Davies P., Fromme B. J., Konvicka K., Chi L., Millar R. P., Davidson J. S., Weinstein H. i Sealfon S. C. (1998) Functional microdomains in G-protein-coupled receptors. The conserved arginine-cage motif in the gonadotropin-releasing hormone receptor. *J. Biol. Chem.* **273**; 10445-19453.

Ballesteros J. A., Jensen A. D., Liapakis G., Rasmussen S. G., Shi L., Gether U. i Javitch J. A. (2001) Activation of the beta 2-adrenergic receptor involves disruption of an ionic lock between the cytoplasmic ends of transmembrane segments 3 and 6. *J. Biol. Chem.* **276**; 29171-29177.

Berson, E. L. (2000) Nutrition and retinal degenerations. *Int. Ophthalmol. Clin.* **40**; 93-111.

Boeckert J. i Pen J. P. (1999) Molecular thinckering of G protein-coupled receptors: an evolutionary success. *The EMBO Journal* **18**; 1723-1729.

Bosch L., Ramon E., Del Valle L. J. i Garriga P. (2003) Structural and functional role of helices I and II in rhodopsin. A novel interplay evidenced by mutations at Gly-51 and Gly-89 in the transmembrane domain. *J. Biol. Chem.* **278**; 20203-20209.

Breit A., Lagace M. i Bouvier M. (2004) Hetero-oligomerization between beta2- and beta3-adrenergic receptors generates a beta-adrenergic signaling unit with distinct functional properties. *J. Biol. Chem.* **279**; 28756-28765.

Bubis J. (1998) Effect of detergents and lipids on transducin photoactivation by rhodopsin. *Biol. Res.* **31**; 59-71.

Buss V., Sugihara M., Entel P. i Hafner J. (2003) Thr94 and Wat2b effect protonation of the retinal chromophore in rhodopsin. *Angew. Chem. Int. Ed Engl.* **42**; 3245-3247.

Cha K., Reeves P. J. i Khorana H. G. (2000) Structure and function in rhodopsin: Destabilization of rhodopsin by the binding of an antibody at the N-terminal segment provides support for involvement of the latter in an intradiscal tertiary structure. *Proc. Natl. Acad. Sci. USA* **97**; 3016-3021.

Chen S., Lin F., Xu M. i Graham R. M. (2002) Phe(303) in TMVI of the $\alpha 1b$ adrenergic receptor is a key residue coupling TM helical movements to G-protein activation. *Biochemistry* **41**; 588-596.

Coddou C., Morales B. i Huidobro-Toro J. P. (2003) Neuromodulator role of zinc and copper during prolonged ATP applications to P2X4 purinoceptors. *Eur. J. Pharmacol.* **472**; 49-56.

Cohen S. N., Chang A. C. Y., i Hsu L. (1972) Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. USA* **69**; 2110-2114 .

Cohen G. B., Yang T., Robinson P. R i Oprian D. D. (1993) Constitutive activation of opsin: influence of charge at position 134 and size at position 296. *Biochemistry* **32**; 6111-6115.

- Courageot M. P., Lepine S., Hours M., Giraud F. i Sulpice J. C. (2004) Involvement of sodium in early phosphatidylserine exposure and phospholipid scrambling induced by P2X7 purinoceptor activation in thymocytes. *J. Biol. Chem.* **279**; 21815-21823.
- Creemers A. F., Kiihne S., Bovee-Geurts P. H., DeGrip W.J., Lugtenburg J. i de Groot H. J. (2002) ¹H and ¹³C MAS NMR evidence for pronounced ligand-protein interactions involving the ionone ring of the retinylidene chromophore in rhodopsin. *Proc. Natl. Acad. Sci. USA* **99**; 9101-9106.
- Cuatrecasas P., Wilcheck M. i Anfinsen C. D. (1968) Selective enzyme purification by affinity chromatography. *Proc. Natl. Acad. Sci. USA* **61**; 636-643.
- Davidson F. F., Loewen P. C. i Khorana H. G. (1994) Structure and function in rhodopsin: replacement by alanine of cysteine residues 110 and 187, components of a conserved disulfide bond in rhodopsin, affects the light-activated metarhodopsin II state. *Proc. Natl. Acad. Sci. USA* **91**; 4029-4033.
- DeGrip, W. J. i Bovee-Geurts P. H. M. (1979) Synthesis and properties of alkylglucosides with mild detergent action: Improved synthesis and purification of 2-1-Octyl-, -Nonyl-, and Decyl-Glucose. Synthesis of 2-1-Undeglycose and 2-1-Dodecylmaltoside. *Chemistry and Physics of Lipids* **23**; 321-335.
- DeGrip W. J., Vanoostrum J. i Bovee-Geurts P. H. (1998) Selective detergent-extraction from mixed detergent/lipid/protein micelles, using cyclodextrin inclusion compounds: a novel generic approach for the preparation of proteoliposomes. *Biochem. J.* **330**; 667-674.
- Del Valle L. J., Escribano C., Perez J. J. i Garriga P. (2002) Calcium-induced decrease of the thermal stability and chaperone activity of alpha-crystallin. *Biochim. Biophys. Acta.* **1601**; 100-109.
- Doi T., Molday R. S. i Khorana H. G. (1990) Role of the intradiscal domain in rhodopsin assembly and function. *Proc. Natl. Acad. Sci. USA* **87**; 4991-4995.
- Dryja T. P., Berson E. L., Rao V. R. i Oprian D. D. (1993) Heterogeneous missense in the rhodopsin gene as a cause of Congenital Stationary Night Blindness. *Nature Genetics* **4**; 280-282.
- Duñach M., Sabés M. i Padrós E. (1983) Fourth derivative spectrophotometry analysis in tryptophan environment in proteins. *Biochemistry* **134**; 17448-17457.
- Eason M. G. i Liggett S. B. (1996) Chimeric Mutagenesis of putative G-protein Coupling Domains of the α_2 -adrenergic receptors. *J. Biol. Chem.* **271**; 12826-12832.

Fahmy K. i Sakmar T. P. (1993) Regulation of the rhodopsin-transducin interaction by a highly conserved carboxylic acid group. *Biochemistry* **32**; 7229-7236.

Fahmy K., Siebert F. i Sakmar T. P. (1995) Photoactivated state of rhodopsin and how it can form. *Biophys. Chem.* **56**; 171-181.

Fahmy K., Sakmar T. P. i Siebert F. (2000) Transducin-dependent protonation of glutamic acid 134 in rhodopsin. *Biochemistry* **39**; 10607-10612.

Farrar G. J., Kennan P. F. i Humphries P. (2002) On the genetics of Retinitis Pigmentosa and on mutation-independent approaches to therapeutic intervention. *The EMBO Journal* **21**; 857-864.

Farrens D. L. i Khorana H. G. (1995) Structure and function in rhodopsin. Measurement of the rate of Metarhodopsin II decay by fluorescence spectroscopy. *J. Biol. Chem.* **270**; 5073-5076.

Farrens D. L., Altenbach C., Yang K., Hubbell W. i Khorana H. G. (1996) Requirement of Rigid-Body Motion of Transmembrane helices for light activation of rhodopsin. *Science* **276**; 768-770.

Feng W. i Song Z. H. (2003) Effects of D3.49A, R3.50A, and A6.34E mutations on ligand binding and activation of the cannabinoid-2 (CB2) receptor. *Biochem. Pharmacol.* **65**; 1077-1085.

Ferretti L., Karnik S. S., Khorana H. G., Nassal M. i Oprian D. D. (1986) Total synthesis of a gene for bovine rhodopsin. *Proc. Natl. Acad. Sci. USA* **83**; 599-603.

Filipek S., Krzysko K. A., Fotiadis D., Liang Y., Saperstein D. A., Engel A. i Palczewski K. (2004) A concept for G protein activation by G protein-coupled receptor dimers: the transducin/rhodopsin interface. *Photochem. Photobiol. Sci.* **3**, 628-638.

Fotiadis D., Liang Y., Filipek S., Saperstein D. A., Engel A. i Palczewski K. (2003) Rhodopsin dimers in native disc membranes. *Nature* **421**; 127-128.

Francesconi A. i Duvoisin R. M. (2004) Divalent cations modulate the activity of metabotropic glutamate receptors. *Neurosci. Res.* **75**; 472-479.

Franke R. R., Sakmar T. P., Oprian D. D. i Khorana H. G. (1988) A single amino acid substitution in Rhodopsin (Lysine 248 → Leucine) prevents activation of transducin. *J. Biol. Chem.* **268**; 2119-2122.

Franke R. R., König B., Sakmar T. P., Khorana H. G. i Hofmann K. P. (1990) Rhodopsin mutants that bind but fail to activate transducin. *Science* **250**; 123-125.

Franke R. R., Sakmar T. P., Graham R. M. i Khorana H. G. (1992) Structure and function in rhodopsin. Studies of the interaction between the rhodopsin cytoplasmic domain and transducin. *J. Biol. Chem.* **267**; 14767-14774.

Fraser C. M., Chung F. Z., Wang C. D. i Venter J. C. (1988) Site-Directed Mutagenesis of Human β -adrenergic receptors. Substitution of Aspartic acid 130 by Asparagina produces a receptor with affinity agonist binding that in unclouped from adenilate cyclasa. *Prog. Natl. Acad. Sci. USA* **85**; 5478-5482.

Futterman S. i Rollins M. H. (1973) The catalytic isomerization of all-trans-retinal to 9-cis-retinal and 13-cis-retinal. *J. Biol. Chem.* **248**; 7773-7779.

Garcia-Segura J. M., Gavilanes J. G., Martínez del Pozo A., Montero F., Oñaderra M. i Vivanco F. (1996) Emisión de fluorescencia. En *Técnicas instrumentales de análisis bioquímica*. Proyecto Editorial Ciencias Químicas. pp 120-166.

Garriga P., Liu X. i Khorana H. G. (1996) Structure and function in rhodopsin: correct folding and misfolding in point mutants at and in proximity to the site of the retinitis pigmentosa mutation Leu-125-->Arg in the transmembrane helix C. *Proc. Natl. Acad. Sci. USA* **93**; 4560-4564.

Garwin G. G. i Saari J. C. (2000) High-performance liquid chromatography analysis of visual cycle retinoids. *Methods Enzymol.* **316**; 313-324.

Gerlach L. O., Jakobsen J. S., Jensen K. P., Rosenkilde M. R., Skerlj R. T., Ryde U., Bridger G.J. i Schwartz T.W. (2003) Metal ion enhanced binding of AMD3100 to Asp262 in the CXCR4 receptor. *Biochemistry* **42**; 710-717.

Gether U., Lin S., Ghanouni P., Ballesteros J. A., Weinstein H. i Kobilka B. K. (1997) Agonists induce conformational changes in transmembrane domains III and VI of the beta2 adrenoceptor. *EMBO J.* **16**; 6737-6747.

Gonzalez A. L. i Joly E. (1995) A simple procedure to increase efficiency of DEAE-dextran transfection of COS cells. *Trends Genet.* **11**; 216-217.

Grahn B. H., Paterson P. G., Gottschall-Pass K.T i Zhang Z. (2001) Zinc and the eye. *J. Am. Coll. Nutr.* **20**; 106-118.

Greasley P. J, Fanelli F., Rossier O., Abuin L. i Cotecchia S. (2002) Mutagenesis and modelling of the alpha(1b)-adrenergic receptor highlight the role of the helix 3/helix 6 interface in receptor activation. *Mol. Pharmacol.* **61**; 1025-1032.

Gross A. K., Rao V. R. i Oprian D. D. (2003) Characterization of rhodopsin congenital night blindness mutant T94I. *Biochemistry* **42**; 2009-2015.

Guzman, Y. (1981) Transformed simian cells support the replication of Early SV40 mutants. *Cell* **23**; 175-182.

Han M., Lin S. W. Minkova M., Smith S. O. i Sakmar T. P. (1996) Functional interaction of transmembrane helices 3 and 6 in rhodopsin. Replacement of phenylalanine 261 by alanine causes reversion of phenotype of a glycine 121 replacement mutant. *J. Biol. Chem.* **271**; 32337-32342.

Han M., Smith S. O. i Sakmar T. P. (1998) Constitutive activation of opsin by mutation of methionine 257 on transmembrane helix 6. *Biochemistry* **37**; 8253-8261.

Han M. i Sakmar T. P. (2000) Assays for activation of recombinant expressed opsins by all-trans-retinals. *Methods Enzymol.* **315**; 251-267.

Hardgrave P. A. (2001) Rhodopsin Structure, Function and Topography. The FriedenWald Lecture. *Invest. Ophthalmol. Vis. Sci.* **42**; 3-9.

Heck M., Schadel S. A., Maretzki D., Bartl F. J., Ritter E., Palczewski K. i Hofmann H. P. (2003) Signalling states of rhodopsin. Formation of the storage form, metarhodopsin III, from active metarhodopsin II. *J. Biol. Chem.* **278**; 3163-3169.

Heck M., Schadel S. A., Maretzki D. i Hofmann K. P. (2003b) Secondary binding sites of retinoids in opsin: characterization and role in regeneration. *Vision Res.* **43**; 3003-3010.

Heckenlively J. R., Boughman J. A. i Friedman L. H. (1988) Pedigree analysis. In *Heckenlively JR. Retinitis Pigmentosa*. Philadelphia PA, Lippincott, pp 6-24.

Herbert T. E., Moffett S., Morello J. P., Loise T.P., Bichet D.G., Barret C., i Bouvier M. (1996) A Peptide from a β_2 -Adrenèrgic Receptor Transmembrane Domain Inhibits Both Receptor Dimerization and Activation. *J. Biol. Chem.* **271**; 16384-16392.

Hessel E., Muller P., Herrmann A. i Hofmann K. P. (2001) Light-induced reorganization of phospholipids in rod disc membranes. *J. Biol. Chem.* **276**; 2538-2543.

Hogger P., Shockley M. S., Lamah J. i Sadee W. (1995) Activating and inactivating mutations in N- and C-terminal i3 loop junctions of muscarinic acetylcholine Hm1 receptors. *J. Biol. Chem.* **270**; 7405-7410.

Holst B. i Schwartz T. W. (2003) Molecular mechanism of agonism and inverse agonism in the melanocortin receptors: Zn(2+) as a structural and functional probe. *Ann. N. Y. Acad. Sci.* **994**; 1-11.

Hong K., Knudsen P. J. i Hubbell W. L. (1982) Purification of rhodopsin on hydroxyapatite columns, detergent exchange, and recombination with phospholipids. *Methods Enzymol.* **81**; 144-150.

Huang P., Li J., Chen C., Visiers I., Weinstein H. i Liu-Chen L. Y. (2001) Functional role of a conserved motif in TM6 of the rat mu opioid receptor: constitutively active and inactive receptors result from substitutions of Thr6.34(279) with Lys and Asp. *Biochemistry* **40**; 13501-13509.

Huang P., Visiers I., Weinstein H. i Liu-Chen L. Y. (2002) The local environment at the cytoplasmic end of TM6 of the mu opioid receptor differs from those of rhodopsin and monoamine receptors: introduction of an ionic lock between the cytoplasmic ends of helices 3 and 6 by a L6.30(275)E mutation inactivates the mu opioid receptor and reduces the constitutive activity of its T6.34(279)K mutant. *Biochemistry* **41**, 11972-11980.

Humphries M. M., Rancourt D., Farrar G. J., Kenna P., Hazel M., Bush R. A., Sieving P. A., Sheils D. M., McNally N., Creighton P., Erven A., Boros A., Gulya K., Capecchi M. R. i Humphries P. (1997) Retinopathy induced in mice by targeted disruption of the rhodopsin gene. *Nat. Genet.* **15**, 216-219.

Hwa J., Garriga P., Liu X. i Khorana H. G. (1997) Structure and function in rhodopsin. Packing of the helices in the transmembrane domain and folding to a tertiary structure in the intradiscal domain are coupled. *Proc. Natl. Acad. Sci. USA* **94**; 10571-10576.

Jäger S., Palczewski K. i Hofmann K. P. (1996) Opsin/all-trans-retinal complex activates transducin by different mechanisms than photolyzed rhodopsin. *Biochemistry* **35**; 2901-2908.

Jansen J. J., Mulder W. R., De Caluwe G. L., Vlaskovits J. M. i DeGrip W. J. (1991) In vitro expression of bovine opsin using recombinant baculovirus: the role of glutamic acid (134) in opsin biosynthesis and glycosylation. *Biochim. Biophys. Acta* **1089**; 68-76.

Janz J. M. i Farrens D. L. (2004) Role of retinal hydrogen bond network in rhodopsin schiff base stability and hydrolysis. *J. Biol. Chem.* Epub October 8; M408766200.

Jastrzebska B., Maeda T., Zhu L., Fotiadis D., Filipek S., Ángel A., Stenkamp R. E. i Palczewski K. (2004) Functional characterization of rhodopsin monomers and dimers in detergent. *J. Biol. Chem.* Epub 15 October; M408691200.

Javitch J. A., Fu D., Liapakis G. i Chen J. (1997) Constitutive activation of the beta2 adrenergic receptor alters the orientation of its sixth membrane-spanning segment. *J. Biol. Chem.* **272**; 18546-18549.

Jensen A. D., Guarnieri F., Rasmussen S. G., Asmar F., Ballesteros J. A. i Gether U. (2001) Agonist-induced conformational changes at the cytoplasmic side of transmembrane segment 6 in the beta 2 adrenergic receptor mapped by site-selective fluorescent labeling. *J. Biol. Chem.* **276**; 9279-9290.

Jobling M. F., Huang X., Stewart L. R., Barnham K. J., Curtain C., Volitakis I., Perugini M., White A. R., Cherny R. A., Masters C. L., Barrow C. J., Collins S. J., Bush A. I. i Cappai R. (2001) Copper and zinc binding modulates the aggregation and neurotoxic properties of the prion peptide PrP106-126. *Biochemistry*. **40**; 8073-8084.

Kalloniatis M. i Fletcher E. L. (2004) Retinitis pigmentosa: understanding the clinical presentation, mechanisms and treatment options. *Clin. Exp. Optom.* **87**; 65-80.

Kaushal S. i Khorana H. G. (1994) Structure and function in rhodopsin. 7. Point mutations associated with autosomal dominant retinitis pigmentosa. *Biochemistry* **33**; 6121-6128.

Kaushal S., Ridge K. D. i Khorana H. G. (1994) Structure and Function in Rhodopsin. The Role of Asparagine-linked glycosylation. *Proc. Natl. Acad. Sci. USA* **91**; 4024-4028.

Kelleher D. J. i Johnson G. L. (1988) Transducin inhibition of light-dependent rhodopsin phosphorylation: evidence for beta gamma subunit interaction with rhodopsin. *Mol. Pharmacol.* **34**; 452-460.

Kim J. M., Altenbach C., Thurmond R. L., Khorana H. G. i Hubbell W. L. (1997) Structure and function in rhodopsin: rhodopsin mutants with a neutral amino acid at E134 have a partially activated conformation in the dark state. *Proc. Natl. Acad. Sci. USA* **94**; 14273-14278.

Kjelsberg M. A., Cotecchia S., Ostrowski J., Caron M. G. i Lefkowitz R. J. (1992) Constitutive activation of the alpha 1B-adrenergic receptor by all amino acid substitutions at a single site. Evidence for a region which constrains receptor activation. *J. Biol. Chem.* **267**; 1430-1433.

Konig B., Arendt A., McDowell J. H., Kahlert M., Hardgrave P. A i Hoffmann K. P. (1989) Three Cytoplasmic loops of Rhodopsin interact with transducin. *Proc. Natl. Acad. Sci. USA* **86**; 6878-6882.

Kosugi S., Mori T. i Shenker A. (1998) An anionic residue at position 564 is important for maintaining the inactive conformation of the human lutropin/choriogonadotropin receptor. *Mol. Pharmacol.* **53**; 894-901.

Kroeger K. M., Pflieger K. D. i Eidne K. A. (2003) G-protein coupled receptor oligomerization in neuroendocrine pathways. *Front. Neuroendocrinol.* **24**; 254-278.

Laemmli U. (1970) Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature* **227**; 680-685.

Lagerstrom M. C., Jovins J., Fredriksson R., Fridmanis D., Haitina T., Ling M. K., Berglund M. M. i Schiöth H. B. (2003) High affinity agonistic metal ion binding sites within the melanocortin 4 receptor illustrate conformational change of transmembrane region 3. *J. Biol. Chem.* **278**; 51521-51526.

Lange R., Frank J., Saldana J. L. i Balny C. (1996) Fourth derivative UV spectroscopy of proteins under high pressure. I. Factors affecting the fourth derivative spectrum of the aromatic amino acids. *Eur. Biophys. J.* **24**, 273-283

Leure-duPree A. E. (1981) Electron-Opaque inclusions in the rat retinal pigment epithelium after treatment with chelators of zinc. *Invest. Ophthalmol. Vis. Sci.* **21**; 1-9.

Leure-duPree A. E. i McClain C. J. (1982) The effect of severe Zinc deficiency on the morphology of the rat retinal pigment epithelium. *Invest. Ophthalmol. Vis. Sci.* **23**; 425-434.

Li J., Huang P., Chen C., de Riel J. K., Weinstein H. i Liu-Chen L. Y. (2001) Constitutive activation of the mu opioid receptor by mutation of D3.49(164), but not D3.32(147): D3.49(164) is critical for stabilization of the inactive form of the receptor and for its expression. *Biochemistry.* **40**; 12039-12050.

Liang Y., Fotiadis D., Filipek S., Saperstein D. A., Palczewski K. i Engel A. (2003) Organization of the G protein-coupled receptors rhodopsin and opsin in native membranes. *J. Biol. Chem.* **278**; 21655-21662.

Liang Y., Fotiadis D., Maeda T., Maeda A., Modzelewska A., Filipek S., Saperstein D.A., Engel A. i Palczewski K. (2004) Rhodopsin signaling and organization in heterozygote rhodopsin knockout mice. *J. Biol. Chem.* **279**; 48189-48196.

Lin S. W. i Sakmar T. P. (1996) Specific tryptophan UV-absorbance changes are probes of the transition of rhodopsin to its active state. *Biochemistry* **35**; 11149-11159.

Lu B., Kwan T., Kurimoto Y., Shatos M., Lund R. D. i Young M. J. (2002) Transplantation of EGF-responsive neurospheres from GFP transgenic mice into the eyes of rd mice. *Brain Res.* **943**, 292-300.

Luecke H., Schobert B., Richter H. T., Cartailler J. P. i Lanyi J. K. (1999) Structure of bacteriorhodopsin at 1.55 Å resolution. *J. Mol. Biol.* **291**; 899-911.

Mandel M. i Higa A. (1970) Calcium dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**, 154-159.

Marin E. P., Krishna G., Zyaga T. A., Isele J., Siebert F. i Sakmar T.P. (2000) The amino terminus of the fourth cytoplasmic loop of rhodopsin modulates rhodopsin-transducin interaction. *J. Biol. Chem.* **275**; 1930-1936.

Marrón J. (1996) Aplicació de tècniques espectroscòpiques per a l'anàlisi funcional del fotorreceptor rodopsina i la caracterització estructural dels seus intermediaris. Tesina dirigida per Joan Manyosa i David Garcia-Quintana. Universitat Autònoma de Barcelona.

Mitchell D. C., Niu S. L. i Litman B. J. (2001) Optimization of receptor-G protein coupling by bilayer lipid composition I: kinetics of rhodopsin-transducin binding. *J. Biol. Chem.* **276**; 42801-42806.

Molday R. S. i MacKenzie D. (1983) Monoclonal antibodies to rhodopsin: characterization, cross-reactivity, and application as structural probes. *Biochemistry* **12**; 4517-4523.

Molday R. S. (1998) Photoreceptor Membrane proteins, phototransduction, and retinal degenerative diseases. *Invest. Ophthalmol. & Vis. Sci.* **39**; 2493-2513.

Nair K. S., Balasubramanian N. i Slepak V. Z. (2002) Signal-dependent translocation of transducin. RGS9-1-Gbeta complex, and arrestin to detergent-resistant membrane rafts photoreceptors. *Curr. Biol.* **12**; 421-425.

Naash M. I., Wu T. H., Chakraborty D., Fliesler S. J., Ding X. Q., Nour M., Peachey N. S., Lem J., Qtaishat N., Al-Ubaidi M. R. i Ripps H. (2004) Retinal abnormalities associated with the G90D mutation in opsin. *J. Comp. Neurol.* **478**; 149-163.

Nathans J. i Hogness D. S. (1984) Isolation and nucleotide sequence of the gene encoding human rhodopsin. *Proc. Natl. Acad. Sci. USA* **81**; 4851-4855.

Nathans J. (1990) Determinants of visual pigment absorbance: identification of the retinylidene Schiff's base counterion in bovine rhodopsin. *Biochemistry* **29**; 9746-9752.

Nicolas C., Ghedira I., Stiemer R., Teufel B., Hecquet C., Faure J. P. i Mirshahi M. (2000) Identification of visual arrestin (S-antigen) in retinal pigmented epithelial cells. *Curr. Eye Res.* **21**; 677-683.

Noorwez S. M., Malhotra R., McDowell J. H., Smith K. A., Krebs M. P. i Kaushal S. (2004) Retinoids assist the cellular folding of the autosomal dominant retinitis pigmentosa opsin mutant P23H. *J. Biol. Chem.* **279**; 16278-16284.

Oishi M. i Cosloy S. D. (1972) The genetic and biochemical basis of the transformability of *Escherichia coli* K12. *Biochem. Biophys. Res. Commun.* **49**; 1568-1572.

Okada T., Fujiyoshi Y., Silow M., Navarro J., Landau E. M i Shichida Y. (2002) Functional role of internal water molecules in rhodopsin revealed by X-ray crystallography. *Proc. Natl. Acad. Sci.* **99**; 5982-5987.

Oprian D. D., Molday R. S., Kaufman R.J. i Khorana H.G. (1987) Expression of a synthetic bovine rhodopsin gene in monkey kidney cells. *Proc. Natl. Acad. Sci. USA* **84**, 8874-8878.

Padrós E., Duñach M., Morros A., Sabés M. i Mañosa J. (1982). The state of tyrosine and phenylalanine residues in proteins analyzed by fourth-derivative spectrophotometry. Histone H1 and ribonuclease A. *Eur. J. Biochem.* **127**; 117-122.

Palczewski K., Kumasaka T., Hori T., Behnke C. A., Motoshima H., Fox B. A., Le Trong I., Teller D. C., Okada T., Stenkamp R. E., Yamamoto M. i Miyano M. (2000) Crystal structure of rhodopsin: A G protein-coupled receptor. *Science*. **289**; 739-745.

Papermaster D. S. i Dreyer W. J. (1974) Rhodopsin Content in the Outer Segment Membranes of Bovine and Frog Retinal Rods. *Biochemistry* **13**; 2438-2444.

Parma J., Duprez L., Van Sande J., Cochaux P., Gervy C., Mockel J., Dumont J. i Vassart G. (1993) Somatic mutations in the thyrotropin receptor gene cause hyperfunctioning thyroid adenomas. *Nature* **365**; 649-651.

Patel A. B., Crocker E., Eilers M., Hirshfeld A., Sheves M. i Smith S. O. (2004) Coupling of retinal isomerization to the activation of rhodopsin. *Proc. Natl. Acad. Sci. USA*. **101**; 10048-10053.

Pauwels P. J., Gouble A. i Wurch T. (1999) Activation of constitutive 5-hydroxytryptamine(1B) receptor by a series of mutations in the BBXXB motif: positioning of the third intracellular loop distal junction and its G(o)alpha protein interactions. *Biochem. J.* **343**; 435-442.

Polozova A. i Litman B. J. (2000) Cholesterol dependent recruitment of di22:6-PC by a G protein-coupled receptor into lateral domains. *Biophys. J.* **79**; 2632-2643.

Pucadyil T. J. i Chattopadhyay A. (2004) Cholesterol modulates ligand binding and G-protein coupling to serotonin(1A) receptors from bovine hippocampus. *Biochim Biophys Acta* **1663**; 188-200.

Ramon E., Marron J., del Valle J. L., Bosch L., Andres A., Manyosa J. i Garriga P. (2003) Effect of dodecil maltoside detergent on rhodopsin stability and function. *Vision Research* **43**; 3055-3061.

Rao R. V., Cohen G. B. i Oprian D. D. (1994) Rhodopsin mutation G90D and a molecular mechanism for congenital night blindness. *Nature* **367**; 639-642.

Rasmussen S. G., Jensen A. D., Liapakis G., Ghanouni P., Javitch J. A. i Gether U. (1999) Mutation of a highly conserved aspartic acid in the beta2 adrenergic receptor: constitutive activation, structural instability, and conformational rearrangement of transmembrane segment 6. *Mol. Pharmacol.* **56**; 175-184.

Ren Q., Kurose H., Lefkowitz R. J. i Cotecchia S. (1993) Constitutively active mutants of the alpha 2-adrenergic receptor. *J. Biol. Chem.* **268**; 16483-16487.

Robinson P. (2000) Assay for detection of constitutively active opsin. *Methods Enzymol.* **315**; 207-218.

Rodriguez J. A., Herrera C. A., Birch D. G. i Daiger S. P. (1993) A leucine to arginine amino acid substitution at codon 46 of rhodopsin is responsible for a severe form of autosomal dominant retinitis pigmentosa. *Hum. Mutat.* **2**; 205-213.

Rosenberg I. M. (1996) Protein Analysis and Purification, Benchtop techniques; Massachusetts General Hospital.

Ruiz A., Borrego S., Marcos I. i Antiñolo G. (2001) Anatomía y Fisiología de la retina. En *La Retinosis Pigmentaria en España*. Coord. Diana Valverde. pp 11-38.

Sagdullaev B. T., Aramant R. B., Seiler M. J., Woch G. i McCall M. A. (2003) Retinal transplantation-induced recovery of retinectal visual function in a rodent model of retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* **44**, 1686-1695.

Sakamoto T. i Khorana H. G. (1995) Structure and function in rhodopsin. The fate of opsin formed upon the decay of light-activated metarhodopsin II *in vitro*. *Proc. Natl. Acad. Sci. USA* **92**; 249-253.

Sakmar T. P., Franke R. R i Khorana H. G. (1989) Glutamic acid-113 serves as the retinylidene Schiff Base Counterion in Bovine Rhodopsin. *Proc. Natl. Acad. Sci. USA* **86**; 8309-8313.

Sakmar T. P. (1998) Rhodopsin: A prototypical G Protein-Coupled Receptor. *Prog. Nucleic Acid Res. Mol. Biol.* **59**;1-34.

Salesse C., Boucher F. i Leblanc R. M. (1984) An evaluation of purity criteria for bovine rod outer segment membranes. *Anal. Biochem.* **142**; 258-266.

Samama P., Cotecchia S., Costa T. i Lefkowitz R. J. (1993) A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model. *J. Biol. Chem.* **268**; 4625-4636.

Schadel S. A., Heck M., Maretzki D., Filipek S., Teller D. C., Palczewski K. i Hofmann K. P. (2003) Ligand channeling within a G-protein-coupled receptor. The entry and exit of retinals in native opsin. *J. Biol. Chem.* **278**; 24896-24903.

Scheer A., Fanelli F., Costa T., De Benedetti P. G. i Cotecchia S. (1996) Constitutively active mutants of the alpha 1B-adrenergic receptor: role of highly conserved polar amino acids in receptor activation. *EMBO J.* **15**; 3566-2578.

Schetz J. A. i Sibley D. R. (1997) Zinc allosterically modulates antagonist binding to cloned D1 and D2 dopamine receptors. *J. Neurochem.* **68**; 1990-1997.

Schetz J. A. i Sibley D. R. (1999) Metal-ions as atomic scale probes of protein structure in *Structure-function analysis of G-protein coupled receptors*. Ed. J. Wess) Wiley-Liss Inc. New York. pp 121-140.

- Shapiro D. A., Kristiansen K., Weiner D. M., Kroeze W. K. i Roth B. L. (2002) Evidence for a model of agonist-induced activation of 5-hydroxytryptamine 2A serotonin receptors that involves the disruption of a strong ionic interaction between helices 3 and 6. *J. Biol. Chem.* **277**; 11441-11449.
- Sheikh S. P., Zyaga T. A., Lichtarge O., Sakmar T. P. i Bourne H. R. (1996) Rhodopsin activation blocked by metal-ion-binding sites linking transmembrane helices C and F. *Nature.* **383**; 347-350.
- Shi L., Liapakis G., Xu R., Guarnieri F. i Ballesteros J. A. (2002) Javitch JA. Beta2 adrenergic receptor activation. Modulation of the proline kink in transmembrane 6 by a rotamer toggle switch. *J. Biol. Chem.* **277**; 40989-40996.
- Shuster T. A., Nagy A. K., Conly D. C. i Farber D. B. (1992) Direct zinc Binding to purified rhodopsin and discs membranes. *Biochem. J.* **282**; 123-128.
- Shuster T. A., Martin F. i Nagy A. K. (1996) Zinc causes an apparent increase in rhodopsin phosphorylation. *Curr. Eye. Res.* **15**; 1019-1024.
- Sieving P. A., Richards J. E., Naarendorf F., Bingham E. L., Scott K. i Alpern M. (1995) Dark-light: Model for nightblindness from de human rhodopsin Gly90—Asp mutations. *Proc. Natl. Acad. Sci. USA* **92**; 880-884.
- Smith W. C., Dinculescu A., Peterson J. J. i McDowell J. H. (2004) The surface of visual arrestin that binds to rhodopsin. *Mol. Vis.* **10**; 392-398.
- Spooner P. J., Sharples J. M., Verhoeven M. A., Lugtenburg J., Glaubitz C. i Watts A. (2002) Relative orientation between the beta-ionone ring and the polyene chain for the chromophore of rhodopsin in native membranes. *Biochemistry.* **41**; 7549-7555.
- Stanasila L., Perez J. B., Vogel H. i Cotecchia S. (2003) Oligomerization of the alpha 1a-and alpha 1b-adrenergic receptor subtypes. Potential implications in receptor internalization. *J. Biol. Chem.* **278**; 40239-40251.
- Stojanovic A., Stitham J. i Hwa J. (2004) Critical role of transmembrane segment zinc binding in the structure and function of rhodopsin. *J. Biol. Chem.* **279**; 35932-35941.
- Strosberg A. D. (1991) Structure/Function Relationship of Protein Belonging to the family of Receptors Coupled to GTP-Binding Protein. *Eur. J. Biochem.* **196**; 1-10.
- Sugihara M., Buss V., Entel P., Elstner M. i Frauenheim T. (2002) 11-cis-retinal protonated Schiff base: influence of the protein environment on the geometry of the rhodopsin chromophore. *Biochemistry.* **41**; 15259-15266.
- Sung C. H., Schneider B. G., Agarwal N., Papermaster D. S. i Nathans J. (1991) Functional heterogeneity of mutant rhodopsins responsible for autosomal dominant retinitis pigmentosa. *Proc. Natl. Acad. Sci. USA* **88**; 8840-8844.

Swaminath G., Lee T. W. i Kobilka B. (2003) Identification of an allosteric binding site for Zn²⁺ on the beta2 adrenergic receptor. *J. Biol. Chem.* **278**; 352-356.

Tam S. W., Wilber K. E. i Wagner F. W. (1976) Light Sensitive Zinc Content of protein fractions from Bovine Rod Outer Segments. *Biochem. Biophys. Res. Commun.* **72**; 302-309.

Teller D. C., Okada T., Behnke C. A. i Palczewski K. (2001) Advances in Determination of a High Resolution Three-Dimensional Structure of Rhodopsin, a Model of g-protein-Coupled Receptors (GPCR's). *Biochemistry* **40**; 7761-7772.

Terakita A., Koyanagi M., Tsukamoto H., Yamashita T., Miyata, T. i Shichida Y. (2004) Counterion displacement in the molecular evolution of the rhodopsin family. *Nat. Struct. Mol. Biol.* **11**; 284-289.

Trumpp-Kallmeyer S., Chini B., Mouillac B., Barberis C., Hoflack J. i Hibert M. (1995) Towards understanding the role of the first extracellular loop for the binding of peptide hormones to G-protein coupled receptors. *Pharm. Acta Helv.* **70**; 255-262.

Vaheri A. i Pagano J. S. (1965) Infectious poliovirus RNA: a sensitive method of assay. *Virology* **27**; 434-436.

Vogel R., Siebert F., Zhang X. Y., Fan G. i Sheves M. (2004). Formation of Meta III during the Decay of Activated Rhodopsin Proceeds via Meta I and Not via Meta II. *Biochemistry* **43**; 9457-9466.

Vorobjev V. S., Sharonova I. N., Sergeeva O. A. i Haas H. L. (2003) Modulation of ATP-induced currents by zinc in acutely isolated hypothalamic neurons of the rat. *Br. J. Pharmacol.* **139**; 919-926.

Wald G. i Brown P. K. (1953) The molar extinction of rhodopsin. *J. Gen. Physiol.* **37**; 189-200.

Wessling-Resnick M. i Johnson G. L. (1987) Allosteric behaviour in transducin activation mediated by rhodopsin. Initial rate analysis of guanine nucleotide exchange. *J. Biol. Chem.* **262**; 3697-3705.

Wray W., Boulikas T., Wray V. P. i Hancock R. (1981) Silver staining of proteins in polyacrylamide gels. *Anal. Biochem.* **118**; 197-203.

Wu S. H., Qiao X., Noebels J. L. i Yang X. L. (1993) Localization and Modulatory actions of zinc in vertebrate retina. *Vision Res.* **33**; 2611-2616.

Yamashita T., Terakita A. i Shichida Y. (2000) Distinct Roles of the Second and Third Cytoplasmic Loops of Bovine Rhodopsin in G-Protein Activation. *J. Biol. Chem.* **275**; 34272-34279.

Yan E. C., Kazmi M. A., De S., Chang B. S., Seibert C., Marin E. P., Mathies R. A. i Sakmar T. P. (2002) Function of extracellular loop 2 in rhodopsin: glutamic acid 181 modulates stability and absorption wavelength of metarhodopsin II. *Biochemistry* **41**; 3620-3627.

Zeng F. Y. i Wess J. (1999) Identification and Molecular Characterization of m3 Muscarinic receptor Dimers. *J. Biol. Chem.* **274**; 19487-19497.

Zimmermann K., Ritter E., Bartl F. J., Hofmann K. P. i Heck M. (2004) Interaction with transducin depletes metarhodopsin III: A regulated retinal storage in visual signal transduction? *J. Biol. Chem.* **279**; 48112-48119.